

Cryp.

edländer
n, Berlin.

ROYAL GARDENS, KEW.

William Marshall

H A N D - L I S T

(906)

OF

HERBS AND FERN ALLIES

CULTIVATED IN

THE ROYAL GARDENS.



L O N D O N :

SOLD AT THE ROYAL GARDENS, KEW.

PRINTED FOR HER MAJESTY'S STATIONERY OFFICE,
BY EYRE AND SPOTTISWOODE,
PRINTERS TO THE QUEEN'S MOST EXCELLENT MAJESTY.

1895.

Price Sixpence.

410276

OK
523
K43X
BOT.

ROYAL / GARDENS, / KEW.



H A N D - L I S T

OF

FERNS AND / FERN ALLIES //

CULTIVATED IN

THE ROYAL GARDENS.



L O N D O N :

SOLD AT THE ROYAL GARDENS, KEW.

PRINTED FOR HER MAJESTY'S STATIONERY OFFICE,
BY EYRE AND SPOTTISWOODE,
PRINTERS TO THE QUEEN'S MOST EXCELLENT MAJESTY.

1895.

Price Sixpence.

ROYAL GARDENS, KEN.



PREFACE

HAND-LIST

VERNS AND TERN ALLES

CONTENTS

THE ROYAL GARDENS



LONDON

PRINTED BY THE ROYAL GARDENS, KEN.
FOR THE ROYAL GARDENS, KEN.
BY THE ROYAL GARDENS, KEN.
PRINTED BY THE ROYAL GARDENS, KEN.

1893

Printed by the Royal Gardens, Ken.

P R E F A C E.

THE general purpose of the series of Hand-lists of the plants cultivated in the Royal Gardens which is in course of publication, and of which the present one is a part, is sufficiently indicated in the preface to that which enumerates the Trees and Shrubs.

The collection of Ferns, whether tropical or temperate, is perhaps, next to that of Palms, the most important feature of the cultivation under glass in the Royal Gardens.

A special interest attaches to it, because it has been used so much in the preparation of most of the books and papers dealing with ferns which have been published in this country during the last half century. The celebrated botanical artist, Francis Bauer, was resident draughtsman for 50 years (1790-1840) to the Royal Botanic Garden at Kew, and the beautiful series of analytical drawings of ferns which he made were published by Sir William Hooker, after his death in 1842, with descriptive letter-press and additions, under the title of *Genera Filicum*. During the 25 years (1840-1865) for which Sir William Hooker was the director of the gardens his attention was specially concentrated on ferns. Under the title of *Species Filicum* he published an elaborate work in five volumes, in which all the known species of typical ferns are fully described, and their synonymy and geographical distribution worked out, with copious illustrations by Mr. Walter Fitch.

Of this, the first volume appeared in 1846, and the last in 1864. He planned also a condensed hand-book of all the known ferns, to be called *Synopsis Filicum*, intended specially for the use of colonists and travellers. All the drawings for this had been made, but the letter-press was not far advanced when he died in the autumn of 1865. It was completed by Mr. J. G. Baker, F.R.S., now Keeper of the Herbarium, in 1868, and a second edition was brought out in 1874. Since that date many hundred new species have been discovered, and a volume of Hooker's *Icones Plantarum*, containing plates and descriptions of the more interesting of the novelties, was published in 1887. Mr. Baker has also published in the *Annals of Botany* (1891) a summary of new ferns discovered or described since 1874. The type specimens from which most of the descriptions and plates contained in this series of books have been made are deposited in the Kew Herbarium.

The living collection in the garden owes its completeness very largely to the zeal and assiduity with which the veteran pteridologist, Mr. John Smith, curator of the Royal Gardens from 1841-63, watched over it for more than 40 years.

In his privately printed *Records of Kew* (pp. 322, 323) he gives the following particulars of its origin and development:—"In the year 1822 I found the collection of ferns at Kew extremely poor, especially as regards tropical species, very many of those introduced in previous years having been lost and very few new ones added. . . . The tender exotic species were more or less of them growing without any arrangement in different houses, and unnamed, their number

amounting to about 40. In 1825 I arranged the tropical species in a group at the end of the then lean-to house . . . now included in the tropical fern house, the area they occupied being 6 ft. by 12 ft. These formed the nucleus of the now great collection. They were successively added to by importations of living plants, as also plants raised from spores obtained from herbarium specimens."

"The collection continued yearly to increase, and in 1846 [1845] I drew up a list of the collection, which was published in an Appendix to the *Botanical Magazine*, for that year; the number then amounted to 400 [378] species. This in 1857 [1856] was followed by another list, entitled 'Catalogue of Cultivated Ferns,' in which 600 [504] species are enumerated."

The principal books published by Mr. Smith are his *Ferns, British and Foreign*, issued in 1866, which contains a classified list of all the species then known in cultivation, full directions for the cultivation of ferns of the different climatic types and by far the most complete history of their gradual introduction which has ever appeared in print, and his *Historia Filicum*, issued in 1875, which contains woodcuts of 220 types and gives a full exposition of his views on fern classification.

In 1868 the last published list was prepared by Mr. J. G. Baker. It enumerates 802 species and varieties of ferns and 48 of fern allies.

The present list comprises 1,116 species and varieties of ferns and 97 of fern allies; this is exclusive of British ferns of which 586 varieties are in cultivation.

The collection falls into three great groups:—(i.) Tropical; (ii.) Temperate; and (iii.) Hardy Ferns.

The TROPICAL FERNS are cultivated in No. II. House.

The TEMPERATE FERNS are cultivated in No. III.; the Filmy Ferns have a separate House (No. II A.) constructed for the special treatment which they require. The fine collection of Temperate Tree ferns is contained in the Temperate House.

The HARDY FERNS are arranged on and about a small rockery, $\frac{7}{E}$ of the Key Plan, and forming the south boundary of Lawn L.

The structural development of the buildings in which the collection is housed has kept pace with its growth. Their history may be given briefly.

The Tropical Fern House (No. II.) is a span-roofed house 129 feet long, 34 feet broad, and 15 feet high in the centre. It has a wide transept on the south side, 40 feet long, 33 feet wide, and 19 feet high.

In 1841 the site of this house was occupied by two lean-to houses used for miscellaneous collections of stove plants. They are enumerated in Dr. Lindley's Report to the Treasury (1838) as Nos. 2 and 3; the former stood to the west and was 50 feet long; the length of the other was 60 feet. It was in No. 2 that, as stated above, John Smith first arranged the nucleus of the collection of Tropical Ferns.

In 1843, No. 2 was doubled and made into a span house.

In 1845, No. 3 was entirely taken down and a new span-roofed house erected in its place. It joined on to

No. 2, which was kept at a somewhat lower temperature, but opened into it through a partition. The centre was filled up with slate shelf staging, and the sides with stone shelves. No. 2 was used as a stove for rare Palms and large Tropical Ferns. No. 3 was at first occupied by the Woburn collection of orchids presented by Her Majesty the Queen; for this purpose it proved eventually unsuited. The orchids were gradually replaced by *Aroidae* and Ferns. Ultimately the latter obtained possession of the whole house. According to John Smith (Records, p. 334), "the centre of the house was a raised sloping rockery, amongst which the plants were growing in a natural state, forming fine bushy specimens."

A small house for tree ferns was built in 1861; this was connected with the principal house in 1868, and now forms the transept.

In 1879 the Tropical Fern House, which from the decay of the timbers had lapsed into an almost ruinous condition, was further severely damaged by the hail storm of August 3rd; 1,152 panes of glass in it were broken. It was subsequently put into a thorough state of repair, but the collections received considerable damage.

In 1887 the west end of the house was set back a few feet; the staging was partly re-arranged and the broad path through the transept was made.

The humid conditions necessary for the cultivation of tropical ferns are unfortunately not conducive to the preservation of the structures in which they are grown when they are built of wood. In 1889 it therefore again

became necessary to renew the east wing of No. II. house. In order to secure greater durability it was decided to use iron for the principal framework, and wood only for the lights. The top-ventilation was provided for by means of a lantern instead of by the older and more clumsy method of sliding sashes. At the same time the use of green glass in the re-constructed wing was abandoned. This was introduced at Kew in 1846 on the recommendation of the late Mr Robert Hunt, F.R.S., who advocated its use on theoretical grounds which are now known to be mistaken. The resulting improvement in the growth of the plants, especially the species of *Adiantum* was very marked, and in 1892 a portion of the west wing was re-glazed with equally satisfactory results.

The TEMPERATE FERNS are now collected in a house (No. III.), which was also built on the system of mixed iron and wood construction. It was erected in 1892 to take the place of a decayed and obsolete structure composed of two small greenhouses united together, Nos. 4 and 5 of Dr. Lindley's Report. The former (the northern wing) was erected in 1803, and in 1840 contained New Holland and Cape plants; the latter, which was "remodelled" in 1825, was filled with succulents.

The reconstructed No. III. is a span-roofed house 60 feet long, 23 feet wide, and 13 feet high. It contains a broad central and two side stages, with paths on both sides.

The north wing of the former No. III. had been latterly devoted almost exclusively to Filmy Ferns after the Kew collection had been enriched by the gift of that formed

by the late John Cooper Forster, which was presented to the Royal Gardens in 1888 by his widow.

The collection is now contained in the new Filmy Fern House (No. II. A.) erected on the north side of No. II. It is 50 feet long by 14 feet wide, with a central path and two cases running the full length of the house.

The HARDY FERNS occupy the rockery originally devoted to Alpine plants (before the construction of the Rock Garden in 1882), and it extends over part of the adjacent ground. A collection was established here in 1874, and the rockery itself was reconstructed in 1888.

The collection as a whole is now probably the richest in existence in garden forms of British species. This is due to the munificent bequest by W. C. Carbonell, Esq., of the extensive collection formed by him at Rhiew Castel, Usk, Monmouthshire. The whole was removed to Kew in 1887. It consisted of 4,261 specimens, many probably unique, besides some hundreds of seedlings.

The total number of well-marked species of ferns and the vascular Cryptogamia (Fern Allies) now mounts up to 3,500. Not more than one-third of these have been brought into cultivation, and Kew is always glad to receive any additions, if possible in the form of well-established plants; if not, in the shape of dry spores, which can be collected and sent very easily by post in small packets.

The following table, which shows the per-centage of the total number of ferns found in different parts of the

world was drawn up by Mr. Baker in 1867; but he does not think that the discoveries of the last 20 years will have altered any of the figures materially :—

Arctic Zone	-	-	-	1 per cent.
Europe	-	-	-	4 „
Temperate Asia, including				
Himalayas	-	-	-	18 „
Temperate N. America			-	5 „
Temperate S. Africa	-	-	-	7 „
Australia and N. Zealand			-	9 „
South temperate America			-	5 „
Tropical Africa	-	-	-	15 „
Tropical Asia	-	-	-	39 „
Tropical America	-	-	-	42 „

The present hand-list is divided into three parts :—


- (i) FERNS proper (pp. 13-133);
- (ii) FERN ALLIES (pp. 135-143); and
- (iii) as an APPENDIX, a list of garden forms of British species (pp. 145-183).

With regard to (i) Ferns proper, the arrangement is alphabetical and synonyms have been only sparingly introduced, those included being chiefly such as are in use in gardens and which are here referred to the correct name in each case. It has not been thought necessary, therefore, to supply an index. Those who wish to pursue the study of fern-nomenclature further may fall back on Hooker and Baker's *Synopsis Filicum*, upon which the present hand-list is substantially based.

The Fern allies (ii) have also been grouped alphabetically. It has not been thought necessary for the

present purpose to separate them according to a botanical classification.

The cultivated and feral varieties of British Ferns (iii) have been enumerated in an Appendix. Although the collection of them at Kew is very rich they stand in a different position to the collection of recognised and well-determined species. Apart from their intrinsic beauty, which is often striking, they are of considerable scientific interest as showing the range of variation due to crossing and seminal reproduction. The amateurs and cultivators who have raised them have furnished them with Latin names, often cumbrous and fantastic, which have received no formal definition. They cannot, therefore, be fixed or quoted for any scientific purpose ; they have, in fact, the same relationship to the species from which they have originated as Bedding Pelargoniums bear to *Pelargonium zonale* or as the Drumhead and other cabbages to *Brassica oleracea*.



I. FERNS.

Acrophorus. *See* Davallia.

Acrostichum, *L.*

A. acuminatum, *Hook.* Brazil.

A. alienum, *Sw.* Tropical America.

A. apodum, *Kaulf.* Tropical America.

A. appendiculatum, *Willd.* Tropical Asia.

A. aureum, *L.* Tropical and sub-tropical regions.

A. Burchellii, *Baker.* Brazil.

A. Cænopteris, *Kze.* Tropical America.

A. cervinum, *Sw.* Tropical America.

A. conforme, *Sw.* Tropical and sub-tropical regions.

Var. *alatum* (*Fée*). Tropical and sub-tropical regions.

A. crassinerve, *Kze.* *See* *latifolium*.

A. crinitum, *L.* Tropical America.

A. crispatulum, *Wall.* *See* *virens*.

A. decoratum, *Kze.* West Indies, Guiana, &c.

A. drynarioides, *Hook.* Penang, Solomon Isles.


A. flagelliferum, *Wall.* Tropical Asia.

A. gramineum, *Baker.* West Indies.

A. Herminieri, *Bory.* West Indies.

A. heteromorphum, *Kl.* Colombia, Ecuador.

A. hybridum, *Bory.* Tropical Africa, Tropical America.



Acrostichum, *L.*—*cont.*

- A. latifolium*, *Sw.* Tropics.
 Var. *angustifolium*, *Hort.* Tropical America.
 Var. *callæfolium* (*Bl.*). Tropical Asia.
 Var. *conopodium*, *Hort.*
 Var. *crassinerve* (*Kze.*). Brazil.
 Var. *Sartorii* (*Liebm.*). Mexico.
- A. Lingua*, *Raddi.* Tropical America.
A. longifolium, *Jacq.* See *latifolium*.
- A. luridum*, *Fée.* Trinidad, Guiana.
- A. magnum*, *Baker.* Demerara.
- A. melanopus*, *Kze.* Venezuela.
- A. muscosum*, *Sw.* Tropical America.
- A. osmundaceum*, *Hook.* Tropical America.
 Var. *frondosum*, *Hort.*
- A. pallidum*, *Baker.* West Indies.
- A. peltatum*, *Sw.* Tropical America.
- A. Prestoni*, *Baker.* Rio Janeiro.
- A. quercifolium*, *Retz.* Tropical Asia.
- A. repandum*, *Bl.* Tropical Asia.
 Var. *quoyanum* (*Gaud.*). Tropical Asia.
- A. Sartorii*, *Liebm.* See *latifolium*.
- A. scandens*, *J. Sm.* Tropical America.
- A. simplex*, *Sw.* Cuba to Brazil.
- A. sorbifolium*, *L.* Tropics.
 Var. *buxifolium* (*Kze.*).
- A. spicatum*, *L.* Tropics of Old World.
- A. squamosum*, *Sw.* Tropics.

Acrostichum, *L.*—*cont.*

- A. Taylora*, *Bailey*. Queensland.
- A. tenuifolium*, *Baker*. Mascarene Islands.
- A. tomentosum*, *Bory*. Bourbon.
- A. undulatum*, *Willd.* *See villosum*.
- A. variabile*, *Hook.* Tropical Asia.
- A. villosum*, *Sw.* Tropical America.
Var. *undulatum* (*Willd.*).
- A. virens*, *Wall.* Tropical Asia.
Var. *crispatulum* (*Wall.*).
- A. viscosum*, *Sw.* Tropics.

Actiniopteris, *Link.*

- A. radiata*, *Link.* Tropical Asia, Tropical Africa.
Var. *australis*, (*Link.*).

Adiantopsis. *See Cheilanthes.**Adiantum*, *L.*

- A. æthiopicum*, *L.* Tropical and south temp. zones.
Var. *assimile* (*Sw.*). Australia.
Var. *chilense* (*Kaulf.*). Chili.
Var. *pellucidum*, (*M. & G.*). Mexico.
Var. *scabrum* (*Kze.*).
- A. affine*, *Hook.* *See diaphanum*.
- A. amabile*, *Moore.* *See Moorei*.
- A. aneitense*, *Carruth.* Aneiteum.
- A. assimile*, *Sw.* *See æthiopicum*.
- A. Bausei*, *Moore.* Garden origin.
- A. bellum*, *Moore.* Bermuda.
- A. Birkenheadii*, *Hort.* Garden origin.

Adiantum, *L.*—*cont.*

- A.* **Capillus-veneris*, *L.* Tropical and temp. zones.
 Var. Lowei, *Hort.* Crimea.
 Var. moritzianum (*Klotzsch*). South America.
- A. cardiochlaena*, *Kze.* *See polyphyllum.*
- A. Catherineæ*, *J. Sm.* *See trapeziforme.*
- A. caudatum*, *L.* Tropics everywhere.
 Var. ciliatum, *Hort.*
- A. Collisii*, *Moore.* Garden origin.
- A. colpodes*, *Moore.* Tropical America.
 Var. elegans, *Hort.*
- A. concinnum*, *H.B.K.* Tropical America.
 Var. Flemingii, *Hort.*
 Var. latum, *Moore.*
- A. cristatum*, *L.* Tropical America.
- A. cultratum*, *J. Sm.* *See trapeziforme.*
- A. cuneatum*, *L. & F.* Tropical America.
 Var. æmulum (*Moore*).
 Var. deflexum, *Hort.*
 Var. dissectum, *Hort.*
 Var. elegans, *Hort.*
 Var. elegantissimum, *Hort.*
 Var. fragrantissimum, *Hort.*
 Var. gracillimum (*Moore*).
 Var. grandis, *Hort.*
 Var. grandiceps, *Hort.*
 Var. lambertianum, *Hort.*
 Var. Lawsoni, *Hort.*

* For garden varieties, *see* p. 145.

Adiantum, *L.*—*cont.*

A. cuneatum, *L. & F.* Tropical America—*cont.*

Var. *Legrandi*, *Hort.*

Var. *mundulum* (*Moore*).

Var. *Pacotti*, *Hort.*

Var. *Rœnbeckii*, *Hort.*

Var. *reginae*, *Hort.*

Var. *schizophyllum*, *Hort.*

Var. *strictum*, *Hort.*

A. curvatum, *Kaulf.* Tropical America.

Var. *brasiliense* (*Raddi*).

A. cyclosorum, *Moore.* Andes.

A. decorum, *Moore.* See *Wagneri*.

A. deflexum. See *cuneatum*.

A. diaphanum, *Bl.* New Zealand, &c.

A. digitatum, *Presl.* Brazil, Peru, &c.

Var. *palmatum* (*Moore*).

Var. *speciosum*, (*Hook.*).

A. dolabriforme, *Hook.* See *lunulatum*.

A. dolosum, *Kze.* Tropical America.

A. Edgeworthii, *Hook.* India and China.

A. elegans, *Hort.* Garden Origin.

A. emarginatum, *Bory.* See *aethiopicum*.

A. excisum, *Kze.* Chili.

Var. *Leyi*, *Hort.*

Var. *multifidum*, *Hort.*

A. Fergusoni, *Moore.* Ceylon.

A. formosum, *R. Br.* Australia.

A. fovearum, *Raddi.* See *tetraphyllum*.

Adiantum, *L.*—*cont.*

- A. fulvum*, *Raoul*. New Zealand.
- A. Funckii*, *Hort.* See *trapeziforme*.
- A. glaucophyllum*, *Hook.* Mexico.
- A. grandiceps*, *Hort.* See *cuneatum*.
- A. Hendersoni*, *Linden.* See *tetraphyllum*.
- A. henslovianum*, *Hook. f.* Colombia, Peru, &c.
- A. hippocrepis*, *Hort.*
- A. hispidulum*, *Sw.* Tropics, &c. of Old World.
 Var. tenellum, *Moore.* Australia.
- A. Hosei*, *Baker.* Borneo.
- A. intermedium*, *Sw.* Tropical America.
- A. Kaulfussii*, *Kze.* Tropical America.
- A. Lathomii*, *Hort.* See *tenerum*.
- A. Legrandi*, *Hort.* See *cuneatum*.
- A. luddemannianum*, *Hort.* Garden origin.
- A. lunulatum*, *Burm.* Tropics everywhere.
 Var. dolabriforme (*Hook.*).
- A. macrophyllum*, *Sw.* Tropical America.
 Var. bipinnatum, *Jenm.* Jamaica.
 Var. albstriatum, *Hort.*
- A. magnificum*, *Hort.* See *Wagneri*.
- A. matthewsianum*, *Hook.* See *polyphyllum*.
- A. monochlamys*, *Eaton.* Japan.
- A. Moorei*, *Baker.* Andes of Peru.
- A. moritzianum*, *Klotzsch.* See *Capillus-veneris*.
- A. neo-caledoniæ*, *Keys.* New Caledonia.
- A. neoguineense*, *Moore.* New Guinea.

Adiantum, *L.*—*cont.*

A. obliquum, *Willd.* Tropical America.

Var. *minus*, *Hort.*

A. Pacotti, *Hort.* See *cuneatum*.

A. pellucidum. See *æthiopicum*.

A. pedatum, *L.* North America, &c.

A. peruvianum, *Klotzsch.* Peru.

A. polyphyllum, *Willd.* Tropical America.

A. prionophyllum, *H.B.K.* See *tetraphyllum*.

A. pulverulentum, *L.* Tropical America.

A. reginæ, *Hort.* See *Capillus-veneris*.

A. reniforme, *L.* Madeira and Teneriffe.

Var. *asarifolium* (*Willd.*). Mauritius.

A. rhodophyllum, *Moore.* Garden origin.

A. rhomboideum, *Hort.* See *obtusum*.

A. Roebelini, *Hort.*

A. rubellum, *Moore.* Bolivia.

A. scabrum, *Kze.* See *æthiopicum*.

A. scutum, *Hort.* See *tenerum*.

A. Seemanni, *Hook.* Guatemala.

A. sessilifolium, *Hook.* See *henslovianum*.

A. setulosum, *J. Sm.* See *diaphanum*.

A. speciosum, *Hook.* See *digitatum*.

A. tenerum, *Sw.* Tropical America.


Var. *alcicorne*, *Hort.*

Var. *Farleyense*, *Moore.*

Var. *Lathomii*, *Hort.*

Var. *scutum*, *Hort.*

Var. *Victoriæ* (*Moore*).



Adiantum, *L.*—*cont.*

A. tetragonum, *Schrad.* Brazil.

A. tetraphyllum, *Willd.* Tropical America.

Var. *acuminatum*, *Hort.*

Var. *gracile*, *Hort.*

A. tinctum, *Moore.* Tropical America.

A. trapeziforme, *L.* Tropical America.

Var. *Catherinæ*, *J. Sm.*

Var. *cultratum*, *J. Sm.*

Var. *Funckii*, *Hort.*

A. triangulatum, *Kaulf.* See *intermedium*.

A. veitchianum, *Moore.* Andes of Peru.

A. Veitchii, *Hance.* See *monochlamys*.

A. velutinum, *Moore.* Tropical America.

A. venustum, *Don.* Himalaya.

Var. *integrifolium*, *Baker.* Himalaya.

A. Victorice, *Moore.* See *tenerum*.

A. villosum, *L.* Tropical America.

A. Wagneri, *Mett.* Andes of Peru.

Var. *magnificum* (*Hort.*).

A. Weigandii, *Moore.* Garden origin.

A. Williamsii, *Moore.* Andes.

A. Wilsoni, *Hook.* See *dolosum*.


Aglaomorpha. See *Polypodium*.

Aleuritopteris. See *Cheilanthes*.

Allosorus. See *Pellæa* and *Cryptogramme*.

Alsophila, *R. Br.*

A. aculeata, *J. Sm.* See *ferox*.



Alsophila, *R. Br.*—*cont.*

- A. Andersoni*, *Scott.* Sikkim.
- A. armata*, *Presl.* Tropical America.
- A. aspera*, *R. Br.* West Indies.
- A. atrovirens*, *Presl.* Tropical America.
- A. australis*, *R. Br.* Australia, &c.
- A. Colensoi*, *Hook. fil.* New Zealand.
- A. contaminens*, *Wall.* Java and Malaya.
- A. crinita*, *Hook.* Ceylon, &c.
- A. excelsa*, *R. Br.* Norfolk Island.
- A. ferox*, *Presl.* Tropical America.
- A. glabra*, *Hook.* East Indies.
- A. infesta*, *Kze.* Andes and Brazil.
 Var. phalerata, *Mart.*
- A. latebrosa*, *Hook.* East Indies.
- A. leichardtiana*, *F. Muell.* Australia.
- A. Malzinei*, *Hort.* Mexico.
- A. paleolata*, *Mart.* Colombia and Brazil.
- A. phalerata*, *Mart.* *See infesta.*
- A. podophylla*, *Hook.* Hong Kong.
- A. procera*, *Kaulf.* Tropical America.
- A. pruinata*, *Kaulf.* Tropical America.
- A. radens*, *Kaulf.* Brazil.
- A. Rebeccæ*, *F. Muell.* Rockingham Bay.
- A. Taenitis*, *Hook.* Brazil.
- A. tomentosa*, *Hook.* East Indies.

Alsophila, *R. Br.*—*cont.*

A. truncata, *Brack.* Fiji, &c.

A. Van Geertii, *Hort.* See *Cyathea mexicana*.

Amphicosmia. See *Hemitelia*.

Anapeltis. See *Polypodium*.

Anchistea. See *Woodwardia*.

Anemia, *Swartz.*

A. collina, *Raddi.* Brazil.

A. dregeana, *Kze.* Natal.

A. fraxinifolia, *Raddi.* See *Phyllitidis*.

A. mandioccana, *Raddi.* Brazil.

A. Phyllitidis, *Sw.* Tropical America.

Var. *foetida*, *Hort.*

A. tomentosa, *Sw.* Tropical America.

Var. *fulva*, (*Sw.*).

Anemidictyon. See *Anemia*.

Anetium. See *Hemionitis*.

Angiopteris, *Hoffm.*

A. evecta, *Hoffm.* Tropics of Old World.

Var. *brongniartiana* (*De Vriese*). Tahiti.

Var. *hypoleuca* (*De Vriese*). Java.

Var. *macrophylla* (*De Vriese*). Java.

Var. *miqueliana* (*De Vriese*). Java.

Var. *pruinosa* (*Kze.*). Java.

Var. *teysmanniana* (*De Vriese*). Java.

Anisogonium. See *Asplenium*.

Antigramme. See *Scolopendrium*.

Arthropteris. See *Polypodium tenellum*, *Nephrodium albo-punctatum*, and *Nephrolepis*.

Aspidium, Swartz.

A. acrostichoides, Sw. North America.

Var. *incisum (Hort.)*.

*A. *aculeatum, Sw.* Ubiquitous.

Var. *angulare (Willd.)*.

Var. *lobatum (Sw.)*.

Var. *moritzianum, (Kze.)*. South America.

Var. *setosum, (Wall.)*. Himalaya.

Var. *tsus-simense (Hook.)*.

A. amabile, Bl. Japan, &c.

A. angulare, Willd. See *aculeatum*.

A. anomalum, H. & A. Ceylon.

A. aristatum, Sw. Tropical Asia.

Var. *coniifolium (Wall.)*.

Var. *variegatum, Hort.*

A. atratum, Wall. See *Nephrodium hirtipes*.

A. auriculatum, Sw. East Indies.

A. Blumei. See *Nephrodium*.

A. capense, Willd. S. Temp. zone, &c.

A. caryotideum, Wall. See *falcatum*.

A. concavum, Hort. See *laserpitiifolium*.

A. confluens. See *Nephrodium*.

A. coniifolium, Wall. See *aristatum*.

A. erythrosorum, Hort. See *Nephrodium*.

A. Fadyeni. See *Fadyenia prolifera*.

A. falcatum, Sw. Japan, &c.

Var. *caryotideum (Wall.)*. East Indies, &c.

* For garden varieties, see p. 145.

Aspidium, Swartz—cont.

A. falcatum, Sw. Japan, &c.—*cont.*

Var. *Fortunei (J. Sm.)*. Japan.

Var. *pendulum, Hort.*

A. falcinellum, Sw. Madeira.

A. flexum, Kze. Juan Fernandez.

A. frondosum, Lowe. Madeira.

A. laserpitiifolium, Mett. Japan.

A. lepidocaulon, Hook. Japan.

A. lobatum, Sw. See *aculeatum*.

A. Lonchitis, Sw. N. temp. zone.

A. macrophyllum, Hort. See *Nephrodium*.

A. marginale. See *Nephrodium*.

A. meniscioides, Schrad. Tropical America.

A. mucronatum, Sw. West Indies.

A. munitum, Kaulf. W. North America.

Var. *imbricans, Hort.*

A. Plumieri, Presl. See *trifoliatum*.

A. proliferum, R. Br. See *aculeatum*.

A. pungens, Kaulf. South Africa.

A. repandum, Willd. Tropical Asia.

A. Richardi, Hook. New Zealand.

A. Standishii, Hort. See *laserpitiifolium*.

A. subtriphyllum, Hook. See *Nephrodium*.

A. triangulum, Sw. West Indies.

Var. *ilicifolium, Fee.*

A. trifoliatum, Sw. Tropical America.

Var. *Plumieri (Presl.)*.

Aspidium, Swartz—cont.

- A. tripterum*, Kze. Japan.
A. uliginosum, Hort. See *Nephrodium cristatum*.
A. variolosum, Wall. See *Nephrodium*.
A. varium, Sw. Japan, &c.
A. vestitum, Sw. See *aculeatum*.

Asplenium, L.

- A. abscissum*, Willd. Tropical America.
A. achilleæfolium, Liebm. Mexico.
A. Adiantum-nigrum, L. North and South temp. zones.
 Var. *acutum* (Willd.).
A. alatum, H. B. K. Tropical America.
A. angustifolium, Mich. North America.
A. anisophyllum, Kze. Natal.
A. apicidens, Moore. See *Vieillardii*.
A. arboreum, Willd. Tropical America.
A. Arnottii, Baker. Sandwich Islands.
A. attenuatum, R. Br. Australia.
 Var. *integrum*, Hort.
A. aureum, Link. See *Ceterach*.
A. auritum, Sw. Tropical America.
 Var. *macilentum* (Kze.).
 Var. *rigidum* (Sw.).
A. australasicum, Hook. See *Nidus*.
A. axillare, Ait. See *umbrosum*.
A. bantamense, Baker. Tropical Asia.
A. Baptistii, Hort. See *Vieillardii*.

Asplenium, *L.*—*cont.*

- A. Belangeri*, *Kze.* Malayan Islands.
- A. brachypterum*, *Kze.* Tropical Africa.
- A. Bradleyi*, *Eaton.* North America.
- A. bulbiferum*, *Forst.* Australia, &c.
 - Var. *fabianum* (*H. & J.*).
 - Var. *laxum* (*R. Br.*).
 - Var. *minus*, *Baker.* New Zealand.
- A. caryotaefolium*, *Hort.* See *cuneatum*.
- A. caudatum*, *Forst.* Tropical and sub-tropical regions.
- A. celtidifolium*, *Kze.* Tropical America.
- A. Ceterach*, *L.* Europe and temp. Asia.
 - Var. *aureum* (*Link*).
- A. cicutarium*, *Sw.* Tropical America.
- A. Colensoi*, *Hook. fil.* New Zealand.
- A. compressum*, *Sw.* St. Helena.
- A. conchatum*, *Moore.* West Indies.
- A. contiguum*, *Kaulf.* Polynesia, &c.
 - Var. *fissum*, *Hort.*
- A. cordifolium*, *Mett.* Malaya, &c.
- A. costale*, *Sw.* West Indies and Peru.
- A. crenulatum*, *Baker.* Tropical America.
- A. cultrifolium*, *L.* West Indies.
- A. cuneatum*, *Lam.* Tropical America, &c.
- A. decurrens*, *Bedd.* India.
- A. decussatum*, *Sw.* Tropics of Old World.
- A. dilatatum*, *Hook.* See *latifolium*.

Asplenium, *L.*—*cont.*

- A. dimidiatum*, *Sw.* Tropical America.
A. dimorphum, *Kze.* Norfolk Island.
A. divaricatum, *Kze.* Chili and Peru.
A. ebenenum, *Ait.* Temp. regions.
A. elongatum, *Sw.* *See tenerum.*
A. erectum, *Bory.* *See lunulatum.*
A. erosum, *L.* West Indies.

A. esculentum, *Presl.* Tropical Asia.
 Var. glabrum, *Hort.*
 Var. serampurens (*Spr.*).

A. fabianum, *H. & J.* *See bulbiferum.*

A. falcatum, *Lam.* Tropics and sub-tropics, Old World.
 Var. erosum, *Hort.*

A. fernandezianum, *Kze.* *See lunulatum.*
A. fijiense, *Brack.* Fiji Islands.
*A. *Filix-fœmina*, *Bernh.* Ubiquitous.
A. firmum, *Kze.* *See abscissum.*
A. flabellifolium, *Cav.* Australia.
A. flaccidum, *Forst.* Australia, &c.
A. fontanum, *Bernh.* North temp. zone.
A. formosum, *Willd.* Tropical America.
A. fragrans, *Sw.* Tropical America.
 Var. fœniculaceum (*H. B. K.*)

* For garden varieties, *see* p. 157.

Asplenium, *L.*—*cont.*

- A. Françonis*, *Mett.* Tropical America.
A. fraxinifolium, *Wall.* See *bantamense*.
A. furcatum, *Thunb.* Tropical and sub-tropical regions.
 Var. *tripinnatum*, *Baker.* Natal.
A. gemmiferum, *Schrad.* South Africa.
A. germanicum, *Weiss.* Europe.
A. Godmani, *Baker.* West Indies.
A. goringianum, *Mett.* See *macrocarpum*.
A. grandifolium, *Sw.* Tropical America.
A. griffithianum, *Hook.* Himalaya.
A. Hemionitis, *L.* South Europe and North Africa.
A. heterocarpum, *Wall.* East Indies.
A. heterophlebium, *Mett.* Himalaya.
A. hians, *Kze.* West Indies.
A. hookerianum, *Colenso.* New Zealand.
A. horridum, *Kaulf.* Sandwich Islands, &c.
A. incisum, *Thb.* Japan.
A. japonicum, *Thb.* Japan.
A. Klotzschii, *Mett.* Colombia and Venezuela.
A. laffanianum, *Baker.* Bermuda.
A. lanceolatum, *Huds.* Europe.
A. lanceum, *Thunb.* Japan.
A. lasiopteris, *Mett.* See *Peterseni*.
A. latifolium, *Don.* Tropical Asia.
A. laxum, *R. Br.* See *bulbiferum*.

THE NEW YORK PUBLIC LIBRARY

ASTOR LENOX TILDEN FOUNDATION

500 FIFTH AVENUE, NEW YORK, N. Y.

Acquired from the

Library of the

City of New York

Gift of the

City of New York

Library of the

City of New York

Gift of the

City of New York

Library of the

City of New York

Gift of the

City of New York

Library of the

City of New York

Gift of the

City of New York

Library of the

City of New York

Gift of the

City of New York

Library of the

City of New York

Gift of the

Asplenium, *L.*—*cont.*

A. lineatum, *Sw.* Mauritius.

Var. *bifidum* (*Presl.*).

A. longissimum, *Bl.* Java, &c.

A. lucidum, *Forst.* See *obtusatum*.

A. lunulatum, *Sw.* Tropics.

Var. *erectum* (*Bory.*).

Var. *lobatum* (*P. & R.*). South Africa.

Var. *pteropus* (*Kaulf.*). Tropical America.

Var. *fernandezianum* (*Kze.*).

Var. *tenellum* (*Roxb.*). St. Helena, &c.

Var. *tripinnatifidum*, *Hort.*

A. macilentum, *Kze.* See *auritum*.

A. macrocarpum, *Bl.* Tropical Asia.

Var. *goringianum* (*Mett.*).

A. macrophyllum, *Sw.* Tropical Asia.

A. marginatum, *L.* Tropical America.

A. marinum, *L.* Europe, &c.

Var. *imbricatum*, *Hort.*

Var. *ramosum*, *Hort.*

A. maximum, *Don.* West Indies to Ceylon.

A. Michauxii, *Spreng.* See *Filix fœmina*.

A. monanthemum, *L.* Temperate regions.

A. monticolum, *Jenm.* Jamaica.

A. Nidus, *L.* Tropics, Old World.

Var. *australasicum*, (*Hook.*). Australia.

A. nigripes, *Bl.* Tropical Asia.

A. nigro-paleaceum, *Baker.* St. Helena.

Asplenium, *L.*—*cont.*

A. niponicum, *Mett.* Japan.

Var. *cristatum*, *Hort.*

A. nitens, *Sw.* Mauritius.

A. nitidum, *Sw.* East Indies.

A. nobile, *Hort.* See *viviparum*.

A. nodulosum, *Hort.* See *lineatum*.

A. novæ-caledoniæ, *Hook.* New Caledonia.

A. obliquum, *Forst.* See *obtusatum*.

A. obtusatum, *Forst.* Australia, &c.

Var. *lucidum* (*Forst.*).

Var. *Lyallii*, *Hook.* New Zealand.

Var. *obliquum* (*Forst.*).

A. obtusifolium, *L.* Tropical America.

A. obtusilobum, *Hook.* Polynesia.

A. oligophyllum, *Kaulf.* Tropical America.

A. paleaceum, *R. Br.* Tropical Australia.

A. paradoxum, *Bl.* Java, &c.

A. Peterseni, *Kze.* East Indies.

Var. *lasiopteris* (*Mett.*).

A. planicaule, *Wall.* Himalaya.

A. plantagineum, *L.* Tropical America.

A. polyodon, *Forst.* See *falcatum*.

A. polypodioides, *Mett.* Tropical Asia.

A. præmorsum, *Sw.* See *furcatum*.

A. proliferum, *Lam.* See *decussatum*.

A. prolongatum, *Hort.* See *rutæfolium*.

A. pteridoides, *Baker.* Lord Howe's Island.

Asplenium, *L.*—*cont.*

- A. pteropus*, Kaulf. *See* *lunulatum*.
A. pubescens, Hort. *See* *esculentum*.
A. reclinatum, Moore. *See* *lunulatum*.
A. refractum, Moore. *See* *fontanum*.
A. resectum, Sm. Tropical Asia.
A. rhizophyllum, Kze. Tropical America.
A. rhizophorum, L. Tropical America.
A. rutæfolium, Kze. Ceylon, South Africa, &c.
 Var. *prolongatum* (*Hook.*).
A. Ruta-muraria, L. North temp. zone.
A. scandens, J. Sm. New Guinea and Philippines.
A. Schimperii, A. Br. East Africa.
A. semihastatum, Kze. Cuba.
A. septentrionale, Hoffm. North temp. zone.
A. Serra, L. & F. Tropical America.
 Var. *natalensis*.
A. serratum, L. Tropical America.
A. Shepherdii, Spreng. Tropical America.
A. sinuatum, Beauv. Tropical Africa.
A. speciosum, Mett. Tropical Asia.
A. splendens, Kze. South Africa.
A. squamulatum, Bl. East Indies.
A. striatum, L. *See* *Shepherdii*.
A. sylvaticum, Presl. Tropics.
A. tenellum, Roxb. *See* *lunulatum*.

Asplenium, *L.*—*cont.*

A. tenerum, *Forst.* Malaya, &c.

Var. *elongatum* (*Sw.*).

A. thelypteroides, *Mich.* North America.

A. Thwaitesii, *A. Br.* Ceylon, &c.

A. tomentosum, *Hook.* East Indies, &c.

*A. *Trichomanes*, *L.* Temp. regions.

A. umbrosum, *J. Sm.* Madeira.

Var. *tenuifolium*, *Kirk.* New Zealand.

A. vagans, *Baker.* Isle of St. Thomas.

A. unilaterale, *Lam.* See *resectum*.

A. veitchianum, *Moore.* See *Belangeri*.

A. Vieillardii, *Mett.* New Caledonia.

Var. *apicidens* (*Moore*).

Var. *Baptistii* (*Hort.*).

A. viride, *Huds.* North temp. zone.

A. viviparum, *Presl.* Mauritius.

Var. *nobile*, *Hort.*

A. zamiioides, *Hook.* See *paradoxum*.

A. zeylanicum, *Hook.* Ceylon.

Athyrium. See *Asplenium*.

Balantium. See *Dicksonia*.

Blechnum, *L.*

B. australe, *L.* South Africa and Madagascar.

B. boreale, *Sw.* See *Lomaria Spicant*.

B. brasiliense, *Desv.* Tropical America.

Var. *corcovadense* (*Raddi*).

* For garden varieties, see p. 169.



Blechnum, Linn.—cont.

- B. cartilagineum*, Sw. Australia.
- B. corcovadense*, Raddi. See *brasiliense*.
- B. Gilliesii*. See *Lomaria procera*.
- B. glandulosum*, Link. See *occidentale*.
- B. gracile*, Kaulf. See *longifolium*.
- B. hastatum*, Kaulf. Temperate South America.
- B. lanceola*, Sw. Tropical America.
- B. latifolium*, Hort. See *longifolium*.
- B. longifolium*, H. B. K. Tropical America.
 Var. *fraxineum* (Willd.).
 Var. *gracile* (Kaulf.).
- B. melanopus*, Hook. India.
- B. occidentale*, L. Tropical America.
 Var. *glandulosum* (Link).
- B. polypodioides*, Raddi. See *unilaterale*.
- B. punctulata*. See *Lomaria*.
- B. rugosum*, Moore. Origin unknown.
- B. triangulare*, Link. See *unilaterale*.
- B. unilaterale*, Willd. Tropical America.
- B. volubile*, Kaulf. Tropical America.

Bolbitis. See *Acrostichum*.

Botrychium, Swartz.

- B. Lunaria*, Sw. Temperate regions.
- B. ternatum*, Sw. North and south temp. zones.
 Var. *dissectum*, Muhl.
- B. virginianum*, Sw. North and south temp. zones.

Brainea, Hook.

B. insignis, *Hook.* Hong Kong and Khasia.

Cænopteris. See *Asplenium*.

Callipteris. See *Asplenium*.

Campyloneuron. See *Polypodium*.

Camptosorus. See *Scolopendrium*.

Campteria. See *Pteris*.

Cassebeera, Kaulf.

C. triphylla, *Kaulf.* South Brazil.

Cephalomanes. See *Trichomanes*.

Ceratopteris, Brong.

C. thalictroides, *Brong.* Tropics.

Ceropteris. See *Gymnogramme*.

Ceterach. See *Asplenium*.

Cheilanthes, Swartz.

C. alabamensis, *Baker.* North America.

C. californica, *Harv.* See *Hypolepis*.

C. chlorophylla, *Sw.* Tropical America.

C. Clevelandii, *Eaton.* California.

C. dicksonioides, *Endl.* See *Hypolepis tenuifolia*.

C. elegans, *Desv.* See *myriophylla*.

C. ellisiana, *Hort.* See *hirta*.

C. farinosa, *Kaulf.* Tropics everywhere.

Var. *anceps* (*Blandf.*). Himalaya.

Var. *punctata*, *Hort.*

C. fragrans, *W. & B.* Europe, &c.

Cheilanthes, Swartz.—cont.

C. hirta, Sw. South America.

Var. *ellisiana*, Hort.

C. multifida, Sw. St. Helena.

C. myriophylla, Desv. Tropical America.

Var. *elegans*, (Desv.).

C. profusa, Kze. See *Pellæa involuta*.

C. pulveracea, Hook. See *Nothochlæna sulphurea*.

C. radiata, R. Br. Tropical America.

C. spectabilis, Kaulf. See *chlorophylla*.

C. tomentosa, Link. North Carolina, &c.

C. vestita, Sw. North America.

Chrysodium. See *Acrostichum*.

Cibotium. See *Dicksonia*.

Cincinalis. See *Nothochlæna*.

Cionidium. See *Deparia*.

Coniogramme. See *Gymnogramme*.

Craspedaria. See *Polypodium*.

Cryptogramme, R. Br.

C. crispa, R. Br. North temp. zone.

Cyathea, Smith.

C. arborea, J. Sm. Tropical America.

C. beyrichiana. See *Hemitelia setosa*.

C. dealbata, Sw. New Zealand.

Var. *tricolor* (Col.).

C. Dregei, Kze. Tropical Africa.

Var. *Burkei*, Hook. South Africa.

THE UNIVERSITY OF CHICAGO

LIBRARY

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

1900

Cyathea, Smith—cont.

C. Hookeri, *Thw.* Ceylon.

C. imrayana, *Hook.* See *Tussacii*.

C. insignis, *Eaton.* West Indies.

C. medullaris, *Sw.* New Zealand.

C. mexicana, *Schlecht.* Mexico.

C. princeps, *J. Sm.* See *insignis*.

C. Schanschin, *Mart.* South America.

C. serra, *Willd.* Tropical America.

C. Smithii, *Hook.* See *Hemitelia Smithii*.

C. spinulosa, *Wall.* India.

C. tricolor, *Col.* See *dealbata*.

C. Tussacii, *Desv.* West Indies.

C. sinuata, *H. & G.* Ceylon.

Cyclodium. See *Aspidium*.

Cyclopeltis. See *Aspidium*.

Cyrtogonium. See *Acrostichum*.

Cyrtomium. See *Aspidium*.

Cyrtophlebium. See *Polypodium*.

Cystopteris, Bernh.

C. alpina, *Desv.* Europe.

C. bergiana. See *Hypolepis*.

C. bulbifera, *Bernh.* North America.

C. fragilis, *Bernh.* Cosmopolitan.

Var. *angustata* (*Sm.*).

Var. *canariensis* (*Moore*).

Var. *dickeana* (*Moore*).

Var. *sempervirens* (*Moore*). Madeira.

Cystopteris, Bernh.—cont.*C. montana*, Link. Europe.*C. tenuis*, Schott. See *fragilis*.*Darea*. See *Asplenium*.*Davallia*, Smith.*D. aculeata*, Sw. West Indies.*D. affinis*, Hook. Tropical and sub-tropical Asia.*D. alpina*, Bl. Malaya, &c.*D. angustata*, Wall. Tropical Asia.*D. assamica*, Baker. Bhotan.*D. bullata*, Wall. Tropical Asia.Var. *decora* (Moore). Java.Var. *Mariesii* (Moore). Japan.*D. canariensis*, Sm. Europe and Canaries.Var. *alopecuroides*.*D. chærophylla*, Wall. East Indies.*D. ciliata*, Hook. Philippines.*D. concinna*, Schr. Tropical Africa and America.*D. cristata*, J. Sm. See *strigosa*.*D. decorata*, Moore. See *bullata*.*D. dissecta*, J. Sm. Java.*D. divaricata*, Bl. Malaya.*D. elata*, Sw. See *elegans*.*D. elegans*, Sw. Tropical Asia, &c.Var. *elata* (Sw.).Var. *Kunzei*, Hort. Malaya.Var. *polydactyla*, Hort.*D. epiphylla*, Bl. Java.

Davallia, Smith--cont.

D. ferruginea, Desv. Madagascar.

D. fijiensis, Hook. Fiji Islands.

Var. *major*, Hort.

Var. *plumosa*, Hort.

D. foeniculacea, Hook. Fiji.

D. griffithiana, Hook. Himalaya, &c.

D. hemiptera, Bory. See *repens*.

D. heterophylla, Sm. Tropical Asia.

D. hirsuta, Sw. Japan, &c.

Var. *angustata*.

D. hirta, Kaulf. Tropical and sub-tropical Asia.

Var. *cristata*.

D. immersa, Wall. Tropical and sub-tropical Asia.

D. khasyana, Hook. See *strigosa*.

D. kunzeana, Hook. See *davallioides*.

D. Lindleyi, Hook. See *solida*.

D. Lorrainei, Hance. Malaya.

D. lonchitidea, Wall. See *platyphylla*.

D. lucida, Hort. See *solida*.

D. marginalis, Baker. Tropical Asia.

D. Mariesii, Moore. See *bullata*.

D. membranulosa, Wall. East Indies.

D. mooreana, Masters. See *pallida*.

D. multidentata, Hk. Himalaya.

D. Novæ zealandiæ, Col. New Zealand.

D. ornata, Wall. See *solida*.

Davallia, *Smith*—*cont.*

- D. pallida*, *Mett.* Malaya and Polynesia.
- D. parvula*, *Wall.* Malaya.
- D. pedata*, *Sm.* Tropical and sub-tropical Asia.
- D. pentaphylla*, *Bl.* Malaya.
- D. pinnata*, *Cav.* East Indies.
- D. platyphylla*, *Don.* East Indies.
- D. polyantha*, *Hook.* *See* *divaricata*.
- D. polypodioides*, *Don.* *See* *Speluncæ*.
- D. pulchra*, *Don.* Himalaya.
- D. pycnocarpa*, *Brack.* *See* *pentaphylla*.
- D. pyxidata*, *Cav.* Australia.
- D. repens*, *Desv.* East Indies.
- D. retusa*, *Cav.* *See* *Lindsaya retusa*.
- D. rhomboidea*, *Wall.* Tropical and sub-tropical Asia.
- D. rubiginosa*, *Kaulf.* Tropical America.
- D. scabra*, *Don.* *See* *marginalis*.
- D. solida*, *Sw.* Tropical Asia and Africa.
 Var. *Lindleyi* (*Hook.*).
 Var. *ornata* (*Wall.*).
- D. Speluncæ*, *Baker.* Tropics, ubiquitous.
- D. strigosa*, *Sw.* Tropical and sub-tropical Asia.
 Var. *rhomboidea* (*Wall.*). North India.
- D. tenuifolia*, *Sw.* Tropics of Old World.
 Var. *stricta*, *Hort.*
 Var. *Veitchii*, *Hort.*
- D. trichosticha*, *Hook.* *See* *Speluncæ*.

Davallia, Smith—cont.

D. Tyermanni, *Baker*. China.

D. vestita, *Bl.* Java and Ceylon.

D. villosa, *Wall.* See *marginalis*.

Dennstædtia. See *Dicksonia*.

Diacalpe, Blume.

D. aspidioides, *Bl.* Tropical Asia.

D. fœniculacea, *C. B. Cl.* See *Aspidium fœniculaceum*.

Dicksonia, Smith.

D. adiantoides, *H. B. K.* Tropical America.

D. antarctica, *Lab.* Australia.

D. apiifolia, *Hook.* Andes of Ecuador and Peru.

D. Barometz, *Link.* Sub-tropical Asia.

D. berteriana, *Hook.* Juan Fernandez.

D. Billardieri. See *antarctica*.

D. Chamissoi, *Hook. fil.* Sandwich Islands.

D. cicutaria, *Sw.* Tropical America.

D. Culcita, *L'Herit.* Madeira, Azores, and Spain.

D. davallioides, *R. Br.* Australia.

D. Deplanchei, *Vieill.* New Caledonia.

D. fibrosa, *Col.* New Zealand.

D. gracilis, *Col.* New Zealand.

D. Herberti, *Moore.* Queensland.

D. Lathamii, *Moore.* Garden origin.

D. Menziesii, *H. & B.* Sandwich Islands.

D. obtusifolia, *Willd.* See *adiantoides*.

D. princeps, *Hort.* See *Cyathea insignis*.

Dicksonia, Smith—cont.

D. regalis, Baker. Mexico.

D. Schiedei, Baker. Tropical America.

D. sellowiana, Hook. South Brazil.

Var. *karsteniana (Klotzch).*

D. spectabilis, Hort. See *Wendlandi*.

D. squarrosa, Sw. New Zealand.

D. tenera, Hort. See *apiifolia*.

D. Wendlandi, Baker. Guatemala.

Var. *Verschaffeltii, Hort.*

D. Youngiæ, C. Moore. New South Wales.

Dictyogramme. See *Gymnogramme*.

Dictyopteris. See *Polypodium*.

Didymochlæna, Desv.

D. lunulata, Desv. Tropics everywhere.

D. sinuosa, Desv. See *lunulata*.

D. truncatula, J. Sm. See *lunulata*.

Didymoglossum. See *Hymenophyllum*.

Diplazium. See *Asplenium*.

Dipteris. See *Polypodium*.

Disphenia. See *Cyathea*.

Doodia, R. Br.

D. aspera, R. Br. Australia.

Var. *multifida, Hort.*

D. blechnoides, Cun. Australia.

D. caudata, R. Br. Australia and New Zealand.

Var. *scabrida.*

Doodia, R. Br.—cont.

D. dives, *Kze.* Ceylon.

D. kunthiana, *Gaud.* Sandwich Islands.

D. lunulata, *R. Br.* See *media*.

D. media, *R. Br.* Australia and New Zealand.

Doryopteris. See *Pteris*.

Drymoglossum, Presl.

D. carnosum, *Hook.* East Himalaya.

Var. **subcordatum** (*Fee.*). China and Japan.

D. piloselloides, *Presl.* Tropical Asia.

Drynaria. See *Polypodium*.

D. diversifolia, *J. Sm.* See *Polypodium rigidulum*.

D. morbillosa, *J. Sm.* See *P. Heracleum*.

Egenolfia. See *Acrostichum*.

Elaphoglossum. See *Acrostichum*.

Eupodium. See *Marattia*.

Fadyenia, Hook.

F. prolifera, *Hook.* West Indies.

Feea. See *Trichomanes*.

Gleichenia, Smith.

G. circinata, *Sw.* Australia and New Zealand.

Var. **Mendelii**, *Hort.* Australia.

Var. **microphylla** (*R. Br.*).

Var. **semivestita** (*Lab.*).

Var. **Speluncæ** (*R. Br.*).

Gleichenia, *Smith*—cont.

G. dicarpa, *R. Br.* Australia and New Zealand.

Var. *alpina* (*R. Br.*).

Var. *glauca*, *Moore*.

Var. *longipinnata*, *Hort.*

G. dichotoma, *Willd.* Tropical and sub-tropical regions.

Var. *nana*, *Hort.*

G. excelsa, *Sm.* See *longissima*.

G. flabellata, *R. Br.* Australia, &c.

G. longipinnata, *Hort.* See *dicarpa*.

G. longissima, *Bl.* Tropics.

G. rupestris, *R. Br.* Australia.

Var. *gigantea*, *Hort.*

Var. *glaucescens*, *Hort.*

Goniophlebium. See *Polypodium*.

Goniopteris. See *Polypodium*.

Gymnogramme, *Desv.*

G. ampla, *Baker.* Australia.

G. calomelanos, *Kaulf.* Tropical America.

Var. *Alstoni*, *Hort.*

Var. *chrysophylla* (*Kaulf.*).

Var. *cristata*, *Hort.*

Var. *laucheana* (*K. Koch*).

Var. *grandiceps*, *Hort.*

Var. *Martensii* (*Kaulf.*).

Var. *Massoni*, *Hort.*

Var. *peruviana* (*Desv.*).

Var. *Parsonsi*, *Hort.*

G. cantoniensis, *Baker.* China.

Gymnogramme, *Desv.*—*cont.*

- G. caudiformis*, *Hook.* Tropical Asia.
- G. chrysophylla*, *Kaulf.* See *calomelanos*.
- G. decomposita*, *Baker.* Garden origin.
- G. decurrens*, *Hook.* See *elliptica*.
- G. diplazioides*, *Desv.* Tropical America.
- G. dobroydense*, *Hort.* Garden origin.
- G. elegantissima*, *Hort.* See *decomposita*.
- G. elliptica*, *Baker.* Tropical Asia, &c.
- G. Feei*, *Hook.* Malaya.
- G. fraxinea*, *Don.* See *javanica*.
- G. gracilis*, *Hew.* Jamaica.
- G. hamiltoniana*, *Hook.* E. Himalaya.
- G. japonica*, *Hook.* Japan, &c.
- G. leptophylla*, *Desv.* Cosmopolitan.
- G. laucheana*, *K. Koch.* See *calomelanos*.
- G. L'Herminieri*, *Link.* See *calomelanos*.
- G. linkiana*, *Kze.* See *diplazioides*.
- G. Mathewsii*, *Hook.* Peru.
- G. Muelleri*, *Hook.* Australia.
- G. multiceps*, *Hort.* See *calomelanos*.
- G. Parsonsii*. See *calomelanos*.
- G. Pearcei*, *Moore.* Peru.
- Var. *robusta*, *Hort.*
- G. peruviana*, *Desv.* See *calomelanos*.
- G. pulchella*, *Linden.* Tropical America.
- Var. *wettenhalliana*, *Hort.*
- Var. *Mayi*, *Hort.*

Gymnogramme, *Desv.*—*cont.*

G. rupestris, Kze. *See* diplazioides.

G. schizophylla, *Baker.* Jamaica.

Var. *gloriosa*, *Hort.*

G. sulphurea, *Desv.* West Indies.

Var. *grandiceps*, *Hort.*

G. tartarea, *Desv.* Tropical America.

G. tomentosa, *Desv.* South America.

G. Totta, *Schlecht.* Tropics.

G. triangularis, *Kaulf.* Vancouver Island, &c.

G. trifoliata, *Desv.* Tropical America.

G. vestita, *Hk.* N. Himalaya, &c.

Gymnopteris. *See* Acrostichum.

Haplopteris. *See* Vittaria.

Helminthostachys, *Kaulf.*

H. dulcis, *Kaulf.* *See* zeylanica.

H. zeylanica, *Hook.* Tropical Asia, &c.

Hemidictyum. *See* Asplenium.

Hemionitis, *L.*

H. cordata, *Roxb.* East Indies.

H. palmata, *L.* Tropical America.

Hemitelia, *R. Br.*

H. brunoniana, *C. B. Cl.* Himalaya.

H. capensis, *R. Br.* S. Africa and Brazil.

H. horrida *R. Br.* West Indies, &c.

H. setosa, *Mett.* Brazil.

H. Smithii, *Hook.* New Zealand.

H. Walkeræ, *Hook.* Ceylon.

Humata. *See* Davallia.

Hymenodium. *See* Acrostichum.

Hymenolepis. *See* Acrostichum.

Hymenophyllum, *Smith*.

H. abruptum, *Hook*. Jamaica and South America.

H. æruginosum, *Carm*. Isle of Tristan d'Acunha.

H. asplenioides, *Sw*. Tropical America, &c.

H. bivalve, *Sw*. New Zealand.

H. Catherinæ, *Hook*. Jamaica.

H. caudiculatum, *Mart*. Tropical America, &c.

H. chilense, *Hook*. Isle of Chiloe, &c.

H. ciliatum, *Sw*. Tropical America.

H. crispum, *H. B. K*. Tropical America.

H. cruentum, *Cav*. Chili.

H. demissum, *Sw*. Australia and New Zealand.

Var. *nitens*, *Hort*.

H. dilatatum, *Sw*. New Zealand, &c.

Var. *forsterianum*, (*Hort*).).

H. elegans, *Spr*. *See* lineare.

H. elegantulum, *V. D. B*. Andes.

H. flabellatum, *Lab*. Australia and New Zealand.

H. forsterianum, *Hort*. *See* dilatatum.

H. fuciforme, *Sw*. Chili and Juan Fernandez.

H. fucoides, *Sw*. Tropical America.

H. hirsutum, *Sw*. Tropical America.

H. javanicum, *Spreng*. Tropical Asia, &c.

Var. *crispatum* (*Wall*).).

Var. *flexuosum* (*Cunn*).).

Var. *tamariscifolium* (*Hort*).).

Hymenophyllum, *Smith*—cont.

H. lineare, *Sw.* Tropical America.

H. magellanicum, *Willd.* South America.

H. multifidum, *Sw.* New Zealand and Pacific Isles.

H. pectinatum, *Cav.* Chili, Chiloe, &c.

Var. *superbum*, *Hort.*

H. polyanthos, *Sw.* Tropics everywhere.

H. pulcherrimum, *Col.* New Zealand.

H. rarum, *R. Br.* Mauritius, New Zealand, &c.

H. scabrum, *A. Rich.* New Zealand.

H. sericeum, *Sw.* Tropical America.

H. subtilissimum, *Kze.* New Zealand.

H. tamariscifolium, *Hort.* See *javanicum*.

H. tunbridgense, *Sw.* Ubiquitous.

Var. *Wilsoni*, *Hook.*

H. villosum, *Col.* See *polyanthos*.

Hypoderris, *R. Br.*

H. Brownii, *J. Sm.* Trinidad.

Hypolepis, *Bernh.*

H. anthriscifolia, *Presl.* Tropical Africa.

H. bergiana, *Hook.* South Africa.

H. californica, *Hook.* California.

H. distans, *Hook.* New Zealand.

H. millefolia, *Hook.* New Zealand.

H. repens, *Presl.* Tropical America.

H. tenuifolia, *Bernh.* Polynesia, &c.

Isoloma. See *Lindsaya*.

Lastrea. See *Nephrodium*.

L. aristata, Hort. See *Aspidium aristatum*.

Lecanopteris, Bl.

L. carnosa, Bl. Malaya.

Lepicystis. See *Polypodium*.

Leptochilus. See *Acrostichum*.

Leptogramme. See *Gymnogramme*.

Leptopteris. See *Todea*.

Leucostegia. See *Davallia*.

Lindsaya, Dry.

L. davallioides, Bl. Malaya.

L. ensifolia, Sw. Old World Tropics.

L. flabellulata, Dry. Tropical Asia.

L. heterophylla, Dry. Tropical Asia.

L. Lowii. See *Acrostichum sorbifolium*.

L. reniformis, Dry. Guiana.

L. retusa, Mett. Philippines, &c.

Litobrochia. See *Pteris*.

Lomaria, Willd.

L. alpina, Spreng. South temperate zone.

L. attenuata, Willd. Tropical and south temperate regions.

Var. *gigantea* (Kaulf.). South Africa.

Var. *heterophylla* (Hort.).

L. Banksii, Hook. fil. New Zealand.

L. boryana, Willd. Tropical and south temperate regions.

Var. *cycadæfolia* (Colla). Chili.

L. capensis, Willd. See *procera*.

THE UNIVERSITY OF CHICAGO
DIVISION OF THE PHYSICAL SCIENCES

DEPARTMENT OF CHEMISTRY
CHICAGO, ILLINOIS

RECEIVED
JANUARY 10, 1950

TO THE EDITOR OF THE JOURNAL OF CHEMICAL PHYSICS
FROM
DR. J. H. VAN VLECK
CHICAGO, ILLINOIS

Enclosed for the Journal of Chemical Physics are two papers
concerning the theory of the Stark effect in the
hydrogen atom. The first paper is a short note
concerning the Stark effect in the hydrogen atom
in the presence of a magnetic field.

The second paper is a longer paper
concerning the Stark effect in the hydrogen atom
in the presence of a magnetic field.

Very truly yours,
J. H. VAN VLECK

Enclosed for the Journal of Chemical Physics are two papers
concerning the theory of the Stark effect in the
hydrogen atom. The first paper is a short note
concerning the Stark effect in the hydrogen atom
in the presence of a magnetic field.

The second paper is a longer paper
concerning the Stark effect in the hydrogen atom
in the presence of a magnetic field.

Very truly yours,
J. H. VAN VLECK

Enclosed for the Journal of Chemical Physics are two papers
concerning the theory of the Stark effect in the
hydrogen atom. The first paper is a short note
concerning the Stark effect in the hydrogen atom
in the presence of a magnetic field.

Lomaria, Willd.—cont.

- L. ciliata*, Moore. New Caledonia.
- L. crenulata*, Moore. See *Germainii*.
- L. discolor*, Willd. Australia and New Zealand.
 Var. *bipinnatifida*, Hort.
 Var. *falcata* (Spreng).
 Var. *nuda* (R. Br.).
- L. fluviatilis*, Spreng. Australia and New Zealand.
- L. Germainii*, Hook. Chili.
- L. gibba*, Lab. New Caledonia, &c.
 Var. *tincta*, Hort.
- L. gigantea*, Kaulf. See *attenuata*.
- L. Gilliesii*, Hook. & Grev. See *procera*.
- L. heterophylla*, Hort. See *attenuata*.
- L. lanceolata*, Spr. Australia.
- L. L'Herminieri*, Bory. Tropical America.
- L. magellanica*, Desv. See *boryana*.
- L. membranacea*, Col. New Zealand.
- L. minor*, R. Br. See *procera*.
- L. neo-caledonica*, Hort. See *gibba*.
- L. nigra*, Col. New Zealand.
- L. Patersoni*, Spreng. Australia.
 Var. *elongata* (Bl.).
- L. procera*, Spreng. Australia, &c.
- L. pumila*, Raoul. New Zealand.
- L. punctulata*, Kze. South Africa.
- L.* Spicant*, Desv. North temperate zone.

* For garden varieties, see p. 169.

Published Weekly, except on Sundays and Public Holidays

Subscription Price, \$5.00 per Annum in Advance

Single Copies, 15 Cents

Entered as Second-Class Matter, May 2, 1902

Postage Paid at Chicago, Ill.

Acceptance for mailing at Special Rate of Postage provided for in Act of October 3, 1917

Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill.

Editor: J. H. Harkness

Business Manager: J. H. Harkness

Editorial Board: J. H. Harkness, Chairman

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Members of the Editorial Board: J. H. Harkness, J. C. Harkness, J. H. Harkness

Lomaria, Willd.—cont.

L. tuberculata, J. Sm. *See* *procera*.

L. zamifolia, Hort. *See* *boryana*.

Lomariopsis. *See* *Acrostichum*.

L. buxifolium, Fee. *See* *A. sorbifolium*, var. *buxifolium*.

Lonchitis, L.

L. pubescens, Willd. Tropics.

Lopholepis. *See* *Polypodium*.

Lophosoria. *See* *Alsophila*.

Lorinseria. *See* *Woodwardia*.

Lotzea. *See* *Asplenium*.

Lygodietycn. *See* *Lygodium*.

Lygodium, Sw.

L. circinatum, Sw. *See* *dichotomum*.

L. dichotomum, Sw. Tropical Asia.

L. flexuosum, J. Sm. *See* *dichotomum*.

L. Forsteri, J. Sm. *See* *reticulatum*.

L. japonicum, Sw. Tropics, &c., Old World.

L. lanceolatum, Desv. Madagascar.

L. palmatum, Sw. United States.

L. pinnatifidum, Sw. Tropics, &c., Old World.

L. reticulatum, Schk. Polynesia.

L. scandens, Sw. Tropics, &c., Old World.

Var. *microphyllum* (R. Br.).

L. venustum, Sw. West Indies, &c.

L. volubile, Sw. South America.

Marattia, Smith.

M. alata, *J. Sm.* West Indies.

M. attenuata, *Lab.* New Caledonia.

Var. **Moorei** (*Hort.*).

M. Ascensionis, *J. Sm.* See *fraxinea*, var. *purpurascens*.

M. cristata, *J. Sm.* See *fraxinea*, var. *purpurascens*.

M. cicutæfolia, *Kaulf.* Brazil.

M. Cooperi, *Hort.* See *attenuata*.

M. elegans, *Endl.* See *fraxinea*.

M. fraxinea, *Sm.* Tropics, &c., Old World.

Var. **purpurascens** (*De Vriese.*). Ascension Island.

M. Kaulfussii, *J. Sm.* Tropical America.

M. laxa, *Kze.* Mexico.

M. Moorei, *Hort.* See *attenuata*.

M. sorbifolia, *Bory.* See *fraxinea*.

Meniscium, Schreb.

M. angustifolium, *Willd.* West Indies, &c.

M. deltigerum. See *Acrostichum virens*.

M. reticulatum, *Sw.* Tropical America.

Var. **macrophyllum** (*Kze.*).

Var. **oligophyllum**, *Hort.*

M. serratum, *Cav.* Tropical America.

M. simplex, *Hook.* Tropical and sub-tropical Asia.

Mertensia. See *Gleichenia*.

Mesochlæna. See *Didymochlæna*.

Microlepia. See *Davallia*.

Microsorium. See *Polypodium*.

Mohria, Sw.

M. caffrorum, Desv. South Africa.

Var. *achilleæfolia, Hort.*

M. thurifraga. See *caffrorum*.

Monogramme, Sch.

M. seminuda, Baker. West Indies, &c.

Myriopteris. See *Cheilanthes*.

Neottopteris. See *Asplenium*.

Nephrodium, Rich.

N. æmulum, Baker. Britain and Madeira.

N. affine, Lowe. See *Filix-mas*.

N. albopunctatum, Desv. Guinea, Natal, &c.

N. atratum (Wall.). See *hirtipes*.

N. atrovirens (J. Sm.). See *decompositum*.

N. Blumei, Hook. See *intermedium*.

N. boryanum, Baker. Himalaya.

N. calcaratum, Hook. East Indies.

N. canum, Baker. Himalaya.

N. catopterum, Hook. South Africa.

N. chinense, Baker. China and Japan.

N. chrysolobum, Fée. Brazil and Colombia.

N. cicutarium, Baker. Tropics, ubiquitous.

N. confluens, F. Muell. Queensland.

N. coruscum, Hort. See *Aspidium varium*.

N. cristatum, Mich. Europe and North America.

Var. *uliginosum, Newm.*

N. cuspidatum, Baker. West Indies and Ceylon.

Nephrodium, Rich.—cont.

- N. cyatheoides**, *Kaulf.* Tropical Asia.
- N. decompositum**, *R. Br.* Australia and New Zealand.
 Var. glabellum (Cunn.).
 Var. Shepherdii (Kze.).
- N. decurrens**, *Baker.* Tropical Asia.
- N. decursivo-pinnatum**, *Baker.* China and Japan.
- N. deltoideum**, *Desv.* Tropical America.
- N. denticulatum**, *Hook.* Tropical America.
- N. dilatatum*, *Desv.* See *spinulosum*.
- N. dissectum**, *Desv.* Tropics of Old World.
 Var. membranifolium (Presl.).
- N. divergens**, *Willd.* See *effusum*.
- N. effusum**, *Baker.* Tropical America.
 Var. divergens (Willd.).
- N. elongatum*, *H. & G.* See *Filix-mas*.
- N. eriocarpum*, *D.C.* See *odoratum*.
- N. erythrosorum**, *Hook.* Japan.
 Var. prolificum (Maxim.).
- N. extensum**, *Hook.* Ceylon.
- N.* Filix-mas**, *Desv.* Temperate regions, ubiquitous.
 Var. elongatum (Hook. & Gr.). Sub-tropical regions.
 Var. schimperianum (Hochst.). Simla.
- N. floridanum**, *Hook.* Louisiana and Florida.
- N. Fœnisecii*, *Lowe.* See *æmulum*.
- N. glabellum*, *Cunn.* See *decompositum*.

* For garden varieties, see p. 171.

Nephrodium, Rich.—cont.

- N. goldieanum*, *Hook.* North America.
N. Grisebachii, *Baker.* Cuba.
N. hirtipes, *Hook.* Tropical Asia.
N. hispidum, *Hook.* New Zealand.
N. incisum, *Baker.* West Indies.
N. intermedium, *Baker.* Tropical Asia.
N. irregulare, *Brack.* See *latifolium*.
N. Jenmani, *Baker.* Jamaica.
N. lacerum, *Baker.* Japan.
N. latifolium, *Baker.* Polynesia.
N. lepidum, *Moore.* Polynesia.
N. leuzeanum, *Hook.* Tropical Asia.
N. macrophyllum, *Baker.* Tropical America.
 Var. calvatum, *Hort.*
N. marginale, *Mish.* North America.
N. membranifolium, *Presl.* See *dissectum*.
N. molle, *Desv.* Ubiquitous.
 Var. polydactylon, *Hort.* Tropics.
 Var. violascens, *Link.*
N. montanum, *Baker.* Europe.
N. nevadense, *Baker.* North America.
N. noveboracense, *Desv.* North America.
N. odoratum, *Baker.* Tropical Asia, &c.
N. Oreopteris, *Desv.* See *montanum*.
N. patens, *Desv.* Tropics, &c.
N. polymorphum, *Baker.* Tropical Asia.

Nephrodium, Rich.—cont.

N. prolificum, Maxim. *See* erythrosorum.

N. pubescens, Desv. West Indies.

N. recedens, Hook. Ceylon, &c.

N. recurvum, Hort. *See* æmulum.

N. Richardsi, Baker. New Caledonia.

Var. *multifidum*, Hort.

N. rigidum, Desv. North temp. zone.

Var. *argutum* (Kaulf.).

N. rodigasianum, Morr. Samoa.

N. scabrosum, Baker. Nilgherries.

N. setigerum, Baker. Japan and tropical Asia.

Var. *cristatum*, Hort.

N. setosum, Baker. Java.

N. Sieboldii, Hook. Japan.

N. Sloanei, Baker. Tropical America.

N. sparsum, Don. Tropical Asia.

N. spinulosum*, Desv. North temp. zone.

Var. *dilatatum* (Desv.).

N. subtriphyllum, Baker. Tropical Asia.

N. tenericaule, Hook. *See* setigerum.

N. Thelypteris, Desv. North and South temp. zones.

N. Thwaitesii, Baker. Ceylon.

N. Trimeni, Beddome. Ceylon.

N. truncatum, Presl. East Indies, &c.

N. unitum, R. Br. Tropical and sub-tropical regions.

* For garden varieties, *see* p. 175.

Nephrodium, Rich.—cont.

N. varians, Fée. Trinidad.

N. variolosum, Baker. Malaya.

N. villosum, Presl. Tropical America.

Var. *lachnopodium* (*J. Sm.*).

Nephrolepis, Schott.

N. acuta, Presl. Tropics everywhere.

N. Bausei, Hort. Garden origin.

N. biserrata, Schott. See *acuta*.

N. cordifolia, Baker. Tropics everywhere.

Var. *compacta*, Hort.

Var. *pectinata* (*Schott.*).

Var. *philippensis* (*Hort.*).

Var. *tuberosa* (*Presl.*).

N. davallioides, Kze. Java.

Var. *furcans*, Hort.

N. Duffi, Moore. Australia.

N. ensifolia, Presl. See *acuta*.

N. exaltata, Schott. Tropics everywhere.

Var. *Barteri*, Hort. West Tropical Africa.

Var. *hirsutula*, Presl.

N. pectinata, Schott. See *cordifolia*.

N. philippinensis, Hort. See *cordifolia*.

N. Pluma, Moore. Madagascar.

N. ramosa, Moore. Tropics, Old World.

N. recurvata, Hort. See *exaltata*.

N. rufescens, Presl. Tropics.

Var. *tripinnatifida*, Baker. Solomon Islands.

Nephrolepis, Schott.—cont.

N. tuberosa, Presl. *See cordifolia.*

N. undulata, J. Sm. *See cordifolia.*

Neurocallis. *See Acrostichum.*

Niphobolus. *See Polypodium.*

N. rupestris, Spr. *See Polypodium serpens.*

Niphopsis. *See Polypodium.*

Nothochlæna, R. Br.

N. affinis, Hook. Mexico, &c.

N. eckloniana, Kze. South Africa.

N. ferruginea, Hook. South America and West Indies.

N. flavens, Moore. Tropical America.

N. Hookeri, Lowe. *See nivea.*

N. lævis, M. & G. *See sinuata.*

N. lanuginosa, Desv. South Europe, &c.

N. Marantæ, R. Br. South Europe, &c.

N. nivea, Desv. Tropical America.

N. rufa, Presl. *See ferruginea.*

N. sinuata, Kaulf. Tropical America.

N. sulphurea, J. Sm. California, Mexico, &c.

N. trichomanoides, R. Br. Jamaica, Cuba.

Odontosoria. *See Davallia.*

Oleandra, Cav.

O. articulata, Cav. Tropical Africa.

O. nodosa, Presl. Tropical America.

Olfersia. *See Acrostichum.*

Onoclea, L.

O. germanica, Willd. Europe and North America.

O. pennsylvanica, Willd. See *germanica*.

O. sensibilis, L. North America.

Onychium, Kaulf.

O. auratum, Kaulf. Himalaya.

O. japonicum, Kze. Japan and Himalaya.

Var. *multisectum, Henders.*

O. lucidum, Spr. See *japonicum*.

Ophioglossum, L.

O. pendulum, L. Tropics, Old World.

O. vulgatum, L. Ubiquitous.

Osmunda, L.

O. cinnamomea, L. North America, &c.

O. claytoniana, L. North America, &c.

O. interrupta, Mich. See *claytoniana*.

O. japonica, Thunb. See *regalis*.

O. javanica, Bl. Java, &c.

O. regalis, L. North and South temp. regions.

Var. *corymbifera, Hort.*

Var. *cristata, Hort.*

Var. *japonica (Thunb).* Japan, &c.

Var. *palustris, Hort.*

Var. *purpurascens, Hort.*

O. spectabilis, Willd. See *regalis*.

Paragramma. See *Polypodium*.

Pellaea, Link.

P. adiantifolia, Hort. See *hastata*.

Pellæa, *Link.*—*cont.*

- P. alabamensis*, *Baker.* Alabama.
P. andromedæfolia, *Fée.* California and South Africa.
P. atropurpurea, *Link.* North America.
P. brachyptera, *Baker.* California.
P. cordata, *J. Sm.* Mexico.

Var. *flexuosa* (*Link.*).

- P. densa*, *Hook.* North America.
P. falcata, *Fée.* Tropical Asia and Australia.
P. geraniæfolia, *Fée.* Tropics.
P. gracilis, *Hook.* North temp. zone.
P. hastata, *Link.* South Africa.
P. intramarginalis, *J. Sm.* Mexico.
P. involuta, *Baker.* East Africa.
P. mucronata, *Eaton.* California, &c.
P. paradoxa, *Hook.* Australia.
P. ternifolia, *Fée.* Tropical America.
P. wrightiana, *Hook.* See *mucronata*.

Phegopteris. See *Polypodium.*

- P. lachnopoda*, *J. Sm.* See *Nephrodium amplum.*
P. submarginalis, *J. Sm.* See *N. caripense.*
P. trichodes, *J. Sm.* See *N. setigerum.*

Phlebodium. See *Polypodium.**Phymatodes.* See *Polypodium.*

- P. vulgare*, *Presl.* See *Polypodium Phymatodes.*

Platyserium, *Desv.*

- P. æthiopicum*, *Hook.* Tropical Africa.

1894

1. *...* 1894
2. *...* 1894
3. *...* 1894
4. *...* 1894
5. *...* 1894
6. *...* 1894
7. *...* 1894
8. *...* 1894
9. *...* 1894
10. *...* 1894
11. *...* 1894
12. *...* 1894
13. *...* 1894
14. *...* 1894
15. *...* 1894
16. *...* 1894
17. *...* 1894
18. *...* 1894
19. *...* 1894
20. *...* 1894
21. *...* 1894
22. *...* 1894
23. *...* 1894
24. *...* 1894
25. *...* 1894
26. *...* 1894
27. *...* 1894
28. *...* 1894
29. *...* 1894
30. *...* 1894
31. *...* 1894
32. *...* 1894
33. *...* 1894
34. *...* 1894
35. *...* 1894
36. *...* 1894
37. *...* 1894
38. *...* 1894
39. *...* 1894
40. *...* 1894
41. *...* 1894
42. *...* 1894
43. *...* 1894
44. *...* 1894
45. *...* 1894
46. *...* 1894
47. *...* 1894
48. *...* 1894
49. *...* 1894
50. *...* 1894
51. *...* 1894
52. *...* 1894
53. *...* 1894
54. *...* 1894
55. *...* 1894
56. *...* 1894
57. *...* 1894
58. *...* 1894
59. *...* 1894
60. *...* 1894
61. *...* 1894
62. *...* 1894
63. *...* 1894
64. *...* 1894
65. *...* 1894
66. *...* 1894
67. *...* 1894
68. *...* 1894
69. *...* 1894
70. *...* 1894
71. *...* 1894
72. *...* 1894
73. *...* 1894
74. *...* 1894
75. *...* 1894
76. *...* 1894
77. *...* 1894
78. *...* 1894
79. *...* 1894
80. *...* 1894
81. *...* 1894
82. *...* 1894
83. *...* 1894
84. *...* 1894
85. *...* 1894
86. *...* 1894
87. *...* 1894
88. *...* 1894
89. *...* 1894
90. *...* 1894
91. *...* 1894
92. *...* 1894
93. *...* 1894
94. *...* 1894
95. *...* 1894
96. *...* 1894
97. *...* 1894
98. *...* 1894
99. *...* 1894
100. *...* 1894

Platycerium, Desv.—cont.

P. alcicorne, Desv. Australia.

Var. *Hillii (Moore)*.

P. grande, J. Sm. Tropical Asia and North Australia.

P. Stemmaria, Beauv. See *æthiopicum*.

P. Wallickii, Hook. Malaya.

P. Willinckii, Moore. Java.

Platyloma. See *Pellæa*.

Pleocnemia. See *Nephrodium*.

Pleopeltis. See *Polypodium*.

Pleuridium. See *Polypodium*.

Pœcilopteris. See *Acrostichum*.

Polybotrya. See *Acrostichum*.

Polypodium, L.

P. adnascens, Sw. Tropical Asia.

Var. *lævis (J. Sm.)*.

P. adnatum, Kze. South America.

P. albopunctatissimum, Linden. See *crassifolium*.

P. albosquamatum, Bl. Malaya.

P. alpestre, Hoppe. North temp. zone.

Var. *flexile (Newm.)*. Scotland.

P. amœnum, Wall. North India.

P. amphostemon, Kze. See *angustifolium*.

P. angustatum, Sw. Tropical Asia.

P. angustifolium, Sw. Tropical America.

P. appendiculatum, Wall. East Himalaya.

P. areolatum, H. B. K. See *aureum*.

Polypodium, L.—cont.

P. aureum, L. Tropical America.

Var. *areolatum* (H. B. K.).

Var. *glaucum*, Hort.

Var. *pulvinatum* (Link).

Var. *sporodocarpon* (Willd.).

P. Billardieri, R. Br. Australia and New Zealand.

P. biseriale, Baker. Peru, &c.

P. calcareum, Sm. See *Dryopteris*.

P. californicum, Kaulf. California.

P. calvatum, Baker. China.

P. cambricum, L. See *vulgare*.

P. cameroonianum, Hook. Cameroon Mountains.

P. Catherinæ, J. Sm. See *loriceum*.

P. caudiceps, Moore. Formosa.

P. chnoodes, Spreng. Tropical America.

P. clathratum, C. B. Clarke. Simla.

P. colpodes, Kze. See *plesiosorum*.

P. conjugatum, Lam. Tropical Asia.

P. contiguum, Wall. See *longifolium*.

P. coronans, Wall. See *conjugatum*.

P. crassifolium, L. Tropical America.

Var. *albopunctatissimum* (Linden). Brazil.

P. cuspidatum, Bl. See *persicæfolium*.

P. dissimile, Hook. See *chnoodes*.

P. distans, Raddi. See *diversifolium*.

P. diversifolium, Sw. Tropical America.

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

THE HISTORY OF ARTS

Polypodium, L.—cont.

- P. drepanum*, *Hook.* Madeira.
- P. Dryopteris*, *L.* North temp. zone.
 Var. robertianum (*Hoffm.*).
- P. effusum.* *See* *Nephrodium.*
- P. ensifolium*, *Willd.* *See* *angustifolium.*
- P. elasticum*, *Rich.* Tropical America.
- P. fraxinifolium*, *Jacq.* *See* *diversifolium.*
- P. fossum*, *Baker.* Malaya.
- P. Gardneri*, *Mett.* Ceylon.
- P. glaucophyllum*, *Kze.* West Indies.
 Var. glaucum, *Hort.*
 Var. viride, *Hort.*
- P. griseum*, *Liebm.* Mexico and Guatemala.
- P. guatemalense*, *Hook.* Tropical America.
- P. harpeodes*, *Link.* *See* *loriceum.*
- P. hastæfolium*, *Sw.* West Indies.
- P. hastatum*, *Thunb.* Japan, &c.
- P. hemionitideum*, *Wall.* Tropical Asia.
- P. Heracleum*, *Kze.* Tropical Asia.
- P. hexagonopterum*, *Mich.* North America.
- P. hirsutum*, *Sw.* Brazil.
- P. hymenodes*, *Bl.* *See* *superficiale.*
- P. inæquale*, *Moore.* *See* *guatemalense.*
- P. incanum*, *Sw.* South Africa and America.
- P. incurvatum*, *Bl.* Malaya, &c.
- P. intermedium*, *H. & A.* *See* *californicum.*

Polypodium, L.—cont.

P. irioides, *Lam.* Tropics and sub-tropics, Old World.

P. juglandifolium, *Don.* Himalaya.

Var. *tenuicauda* (*Hook.*).

P. karwinskianum, *A. Br.* See *plebeium*.

P. lachnopus, *Wall.* N. India.

P. lævigatum, *Cuv.* Tropical America.

P. lanceolatum, *L.* Tropics.

P. leiorrhizon, *Wall.* Tropical Asia.

P. lepidopteris, *Kze.* Tropical America.

Var. *sepultum* (*Kaulf.*).

P. Lingua, *Sw.* Tropical Asia, &c.

Var. *corymbiferum*, *Hort.*

Var. *Heteractis* (*Mett.*). Himalaya.

P. longissimum, *Bl.* Tropical Asia.

P. longifolium, *Mett.* Malaya.

P. loriceum, *L.* Tropical America.

Var. *pinnatifidum*, *Baker.*

Var. *vacillans* (*Link.*).

P. lycopodioides, *L.* Tropics everywhere.

Var. *maximum*, *Hort.*

Var. *owariense* (*Desv.*). Tropical Africa.

Var. *salicifolium* (*Willd.*). Tropical America.

Var. *squamulosum*, *Kaulf.* Brazil.

P. macrourum, *Baker.* Australia.

P. megalodus, *Schk.* See *tetragonum*.

P. membranaceum, *Don.* N. India.

P. menisciifolium, *L. & F.* Brazil,

Polypodium, *L.*—*cont.*

- P. meyenianum*, *Schott.* Philippines.
P. musæfolium, *Bl.* Malaya.
P. multilineatum, *Wall.* N. India.
P. neriifolium, *Schk.* Tropical America.
P. nigrescens, *Bl.* Malaya, &c.
P. nigripes, *Hook.* Venezuela.
P. nitidum, *J. Sm.* See lycopodioides.
P. owariense, *Desv.* See lycopodioides.
P. pectinatum, *L.* Tropical America.
 Var. Paradiseæ (*L. & F.*).
 Var. plumosum, *Hort.*
P. peltideum, *Link.* See Phymatodes.
P. pennigerum, *Forst.* New Zealand.
P. percussum, *Cav.* Tropical America.
P. persicæfolium, *Desv.* Tropical Asia.
 Var. grandidens, *Kze.* Java.
P. Phegopteris, *L.* North temp. zone.
 Var. cristatum, *Hort.*
P. Phyllitidis, *L.* Tropical America.
 Var. brevifolium (*Link.*).
 Var. obscurum (*Mett.*).
P. Phymatodes, *L.* Tropics, Asia and Africa.
 Var. longipes (*Link.*).
 Var. terminale (*Link.*).
P. pictum, *Baker.* Polynesia.
P. piloselloides, *L.* Tropical America.
P. plebeium, *Schlecht.* Mexico.

Polypodium, *L.*—*cont.*

- P. plesiosorum*, *Kze.* Tropical America.
 Var. *appendiculatum* (*Klotzsch*).
P. Plumula, *H. B. K.* See *elasticum*.
P. preslianum, *Link.* See *neriifolium*.
P. propinquum, *Wall.* Tropical Asia.
P. pulvinatum, *Link.* See *aureum*.
P. pustulatum, *Forst.* Australia and New Zealand.
P. quercifolium, *L.* Tropical Asia, &c.
 . repens, *L.* Tropical America.
 Var. *undulatum*, *Hort.*
P. reptans, *Sw.* Tropical America.
 Var. *asplenioides*, (*Sw.*).
P. rigidulum, *Sw.* N. Australia, &c.
P. robertianum, *Hoffm.* See *Dryopteris*.
P. rostratum, *Hook.* N. India.
P. rubidum, *Hook.* Philippines.
P. rupestre, *Bl.* Java, &c.
P. Scouleri, *H. & G.* N.W. America.
P. sepultum, *Kaulf.* See *lepidopteris*.
P. serpens, *Forst.* Australia.
P. simplex, *Sw.* See *lineare*.
P. spectrum, *Kaulf.* Sandwich Isles.
P. sphærocephalum, *Wall.* See *angustatum*.
P. sporodocarpon, *Willd.* See *aureum*.
P. squamulosum, *Kaulf.* See *lycopodioides*.
P. stigmaticum, *Presl.* Tropical America.

Polypodium, *L.*—*cont.*

P. subauriculatum, *Bl.* Tropical Asia &c.

Var. *Reinwardtii* (*Kze.*).

P. subpetiolatum, *Hook.* Central America.

P. superficiale, *Bl.* East Indies.

P. suspensum, *L.* Tropical America.

P. teniosum, *Hort.* See *angustifolium*.

P. tenellum, *Forst.* Australia, &c.

P. tenuicauda, *Hook.* See *juglandifolium*.

P. tetragonum, *Sw.* Australia, &c.

Var. *megalodus* (*Schk.*).

P. tricuspe, *Sw.* Japan.

P. vacillans, *Link.* See *loriceum*.

P. vacciniifolium, *L. and F.* Tropical America.

Var. *album*, *Hort.*

P. venosum, *Desv.* See *stigmaticum*.

P. verrucosum, *Wall.* Malaya.

*P. *vulgare*, *L.* North temp. zone.

P. Xiphias, *Moore.* Polynesia.

P. Zippelii, *Bl.* See *heterocarpum*.

Polystichum. See *Aspidium*.

Prosaptia. See *Davallia*.

Pseudathyrium. See *Polypodium*.

Pteris, L.

P. aquilina, L. Cosmopolitan.

P. arguta, Ait. South Europe and Azores.

* For garden varieties, see p. 175.

Pteris, L.—cont.

P. argyræa, Moore. See *quadriaurita*.

P. Bausei. See *semipinnata*.

P. biaurita, L. Tropics.

Var. *memoralis variegata*.

P. collina, Raddi. See *palmata*.

P. crassipes, Ag. See *gigantea*.

P. crenata, Sw. See *ensifformis*.

P. cretica, L. Tropical and sub-tropical regions.

Var. *albolineata*, Hort.

Var. *cristata*, Hort.

Var. *Mayi*, Hort.

Var. *nobilis*, Hort.

P. elata, Ag. Tropical America.

P. ensiformis, Burm. Tropical Asia.

Var. *Victoriæ*, Hort.

Var. *cristata*, Hort.

P. flabellata, Thunb. South Africa.

P. Ghiesbreghtii, J. Sm. See *laciniata*.

P. gigantea, Willd. West Indies.

P. glaucovirens, Linden. See *quadriaurita*.

P. heterophylla, L. West Indies.

Var. *internata* (Moore).

P. incisa, Thunb. Tropical and sub-tropical regions.

P. laciniata, Willd. Tropical America.

Var. *Ghiesbreghtii*, J. Sm.

P. leptophylla, Sw. Brazil.

Pteris, *L.*—*cont.*

P. longifolia, *L.* Tropical and sub-tropical regions.

Var. *Mariesii*, *Hort.* Japan.

P. ludens, *Wall.* Tropical Asia.

P. moluccana, *Bl.* Malaya, &c.

P. memorialis, *Willd.* See *biaurita*.

P. nobilis, *Hort.* See *palmata*.

P. Ouvrardii, *Hort.* See *serrulata*.

P. palmata, *Willd.* Tropical America, &c.

Var. *collina* (*Raddi*).

Var. *nobilis*, *Hort.*

P. patens, *Hook.* Tropics of Old World.

P. podophylla, *Sw.* Tropical America, &c.

P. quadriaurita, *Retz.* Tropics.

Var. *argyræa* (*Moore*).

Var. *rubricaulis* (*J. Sm.*).

Var. *tricolor* (*Hort.*).

P. robusta, *Hort.* See *aculeata*.

P. sagittifolia, *Raddi.* Brazil.

P. scaberula, *A. Rich.* New Zealand.

P. semipinnata, *L.* Tropical and sub-tropical Asia.

Var. *Bausei*, *Hort.*

P. serrulata, *L. fil.* China and Japan.

Var. *compacta*, *Hort.*

Var. *cristata*, *Hort.*

Var. *densa*, *Hort.*

Var. *gracilis*, *Hort.*

Var. *gloriosa*, *Hort.*

Pteris. *L.*—*cont.*

P. serrulata, *L. fil.* China and Japan—*cont.*

Var. *Leyi*, *Hort.*

Var. *maxima*, *Hort.*

Var. *Ouvrardii*, *Hort.*

Var. *undulata*, *Hort.*

P. straminea, *Mett.* Chili.

P. sulcata, *Link.* See *quadriaurita*.

P. tremula, *R. Br.* Australia.

Var. *elegans*, *Hort.*

Var. *kingiana* (*Endl.*).

Var. *smithiana*, *Hort.*

P. umbrosa, *R. Br.* Australia.

P. undulata, *Moore.* Fiji.

P. Victoriae, *Hort.* See *ensiformis*.

P. wallichiana, *Ag.* Himalaya.

Pteropsis. See *Tænitis*.

Pycnopteris. See *Nephrodium*.

Rhipidopteris. See *Acrostichum*.

Saccoloma. See *Davallia*.

Sadleria, *Kaulf.*

S. cyathæoides, *Kaulf.* Sumatra and Sandwich Islands.

Sagenia. See *Nephrodium*.

Salpichlæna. See *Lomaria*.

Schelleleipsis. See *Polypodium*.

Schizocæna. See *Cyathea*.

Schizoloma. See *Lindsaya*.

Scolopendrium, Sm.

S. brasiliense, Kze. Brazil.

S. Krebsii, Kze. See *Lomaria punctulata*.

*S. *vulgare*, Sm. North temperate zone.

Selliguea. See *Gymnogramme*.

Sitolobium. See *Davallia*.

Soromanes. See *Acrostichum Cænopteris*.

Stegania. See *Lomaria*.

Stenochlæna. See *Acrostichum*.

Stenolomana. See *Davallia*.

Stenosemia. See *Acrostichum*.

Struthiopteris. See *Onoclea*.

Taeniopsis. See *Vittaria*.

Thyrsopteris, Kze.

T. elegans, Kze. Juan Fernandez.

Todea, Willd.

T. africana, Willd. See *barbara*.

T. barbara, Moore. South Africa, Australia, and New Zealand.

T. Fraseri, H. & G. Australia, &c.

Var. *wilkesiana* (Brack.). Polynesia.

T. grandipinnula, Moore. See *Moorei*.

T. hymenophylloides, R. & L. New Zealand.

T. Moorei, Baker. Lord Howe's Island.

T. pellucida, Carm. See *hymenophylloides*.

T. rivularis, Sieb. See *barbara*.

* For garden varieties, see p. 179.

Todea, Willd.—cont.

T. superba, Col. New Zealand.

T. wilkesiana, Brack. See *Fraseri*.

Trichomanes, L.

T. alatum, Sw. Tropical America.

Var. *attenuatum* (Hook.).

T. angustatum, Carm. See *tenerum*.

T. apiifolium, Presl. Polynesia, &c.

T. auriculatum, Bl. Tropical Asia, &c.

T. Bancroftii, H. & G. Tropical America.

T. bipunctatum, Poir. See *Filicula*.

T. Bojeri, H. & G. See *cuspidatum*.

T. boschianum, Sturm. See *radicans*.

T. crispum, L. Tropical America.

T. cuspidatum, Willd. Mauritius.

T. elongatum, Cunn. See *rigidum*.

T. exsectum, Kze. Chili, &c.

T. Filicula, Bory. Tropical Asia and Africa.

T. fœniculaceum, Bory. Mauritius, &c.

T. humile, Forst. Polynesia, &c.

T. javanicum, Bl. Malaya.

T. Kaulfussii, Hook. *fil.* West Indies.

T. luschnathianum, Presl. See *radicans*.

T. maximum, Bl. Malaya and Polynesia.

T. meifolium, Hort. See *apiifolium*.

T. membranaceum, L. Tropical America.

Trichomanes, *L.*—*cont.*

- T. muscoides*, *Sw.* Tropics.
- T. parviflorum*, *Poir.* See *foeniculaceum*.
- T. parvulum*, *Poir.* Tropics of Old World.
- T. Petersii*, *A. Gray.* Alabama.
- T. pinnatum*, *Sw.* Tropical America.
- T. Prieurii*, *Kze.* Tropical America.
- T. pulcherrimum*, *Col.* New Zealand.
- T. pyxidiferum*, *L.* Tropics everywhere.
 Var. olivaceum (*Kze.*). Tropical America.
- T. quercifolium*, *H. & G.* See *pusillum*.
- T. radicans*, *Sw.* Tropical and sub-tropical regions.
 Var. alabamense, *A. Gray.*
 Var. Andrewsii, *Newm.* Ireland.
 Var. boschianum (*Sturm.*).
 Var. crispum, *Hort.*
 Var. dilatatum, *Hort.*
 Var. frondosum (*Fée.*).
 Var. luschnathianum (*Presl.*). Brazil.
- T. reniforme*, *Forst.* New Zealand.
- T. reptans*, *Sw.* Tropical America.
- T. rigidum*, *Sw.* Tropics everywhere.
 Var. elongatum (*Cunn.*). New Zealand.
- T. scandens*, *L.* Central America.
- T. spicatum*, *Hedw.* West Indies.
- T. strictum*, *Menz.* New Zealand.
- T. tenerum*, *Spreng.* Tropical America.
- T. trichoideum*, *Sw.* Tropical America.
- T. venosum*, *R. Br.* Australia.

Vittaria, Smith.

V. elongata, Sw. Tropics of Old World.

V. lineata, Sw. Tropical America.

V. scolopendrina, Thwaites. Ceylon, &c.

V. stipitata, Kze. Colombia to Peru.

Woodsia, R. Br.

W. glabella, R. Br. North temperate zone.

W. ilvensis, R. Br. North temperate zone.

W. oregana, Eat. Rocky Mountains.

W. polystichoides, Eat. Japan.

W. scopulina, Eat. North America.

Woodwardia, Smith.

W. angustifolia, Sm. See *areolata*.

W. orientalis, Sw. Japan, &c.

W. radicans, Sw. North temperate zone.

Var. *Brownii*, Hort.

THE
HISTORY OF THE
CITY OF
NEW-YORK
FROM
THE
FIRST
SETTLEMENT
TO
THE
PRESENT
TIME
BY
JOHN
BUTLER

VOL. II.
CONTAINING
THE
HISTORY OF THE
CITY OF
NEW-YORK
FROM
THE
FIRST
SETTLEMENT
TO
THE
PRESENT
TIME
BY
JOHN
BUTLER

THE
HISTORY OF THE
CITY OF
NEW-YORK
FROM
THE
FIRST
SETTLEMENT
TO
THE
PRESENT
TIME
BY
JOHN
BUTLER

II. FERN ALLIES.

Azolla, *Lam.*

A. caroliniana, *Willd.* Tropical and sub-tropical America.

A. pinnata, *R. Br.* Australia.

Equisetum, *L.*

E. arvense, *L.* North temperate and arctic regions.

E. fluviatile, *Sm.* See *limosum*.

E. giganteum, *L.* Tropical America.

E. hyemale, *L.* North temperate and arctic regions.

E. limosum, *L.* North temperate and arctic regions.

E. maximum, *Lam.* North temperate regions.

E. palustre, *L.* North temperate and arctic regions.

E. pratense, *Ehrh.* North temperate and arctic regions.

E. robustum, *A. Br.* North America.

E. scirpoides, *Michx.* North temperate and arctic regions.

E. silvaticum, *L.* North temperate and arctic regions.

E. Telmateia, *Ehrh.* See *maximum*.

E. variegatum, *Schleich.* North temperate and arctic regions.

Isoetes, *L.*

I. Duriæi, *Bory.* Mediterranean region.

I. lacustris, *L.* Europe and North America.

Var. *Morei*, *Moore.* Ireland.

Lycopodium, *L.*

L. alpinum, *L.* Europe.

L. complanatum, *L.* Europe.

L. clavatum, *L.* Europe.

L. Hippuris, *Desv.* See *squarrosum*.

Lycopodium, L.—cont.

- L. Hookeri*, Wall. See *squarrosum*.
L. nummularifolium, Bl. Malaya, &c.
L. Phlegmaria, L. Tropics.
L. Selago, L. Europe.
L. squarrosum, Forst. Tropical Asia.
L. strictum, Baker. Madagascar.
L. taxifolium, Sw. Tropical America.

Marsilea, L.

- M. Drummondii*, A. Br. Australia.
 Var. *salvatrix* (Hans.).
M. hirsuta, R. Br. Australia.
M. macropus, Englm. Texas.
M. quadrifoliata, L. Europe and Asia.
M. salvatrix, Hans. See *Drummondii*.

Pilularia, L.

- P. globulifera*, L. Europe.

Psilotum, Sw.


- P. triquetrum*, Sw. Tropics.
 Var. *mascarenica*, Hort.
 Var. *nudum* (Griseb.).

Salvinia, Schreb.

- S. natans*, Hoffm. Sub-tropics of Old World.

Selaginella, Spring.

- S. africana*, A. Br. See *Vogelii*.
S. albonitens, Spring. West Indies.
S. albospica, Hort. See *tenophylla*.
S. amœna, Hort. See *caulescens*.
S. apus, Spring. North America.



Selaginella, *Spring*—*cont.*

S. argentea, *Spring*. *See caulescens*.

S. atroviridis, *Spring*. Tropical Asia.

S. bakeriana, *Bailey*. Queensland.

S. Braunii, *Baker*. China.

S. cæsia, *Hort*. *See uncinata*.

S. canaliculata, *Baker*. Tropical Asia.

S. caudata, *Spring*. *See canaliculata*.

S. caulescens, *Spring*. Tropical Asia.

Var. *amœna*, *Hort*.

Var. *minor*, *Hort*.

S. cognata. *See Lobbii*.

S. concinna, *Spring*. Mascaren Isles, &c.

S. conferta, *Moore*. *See canaliculata*.

S. convoluta, *Spring*. Tropical America.

S. cuspidata, *Link*. Tropical America.

Var. *elongata*, *Spring*.

S. denticulata, *Link*. Europe.

S. denticulata, *Hort*. *See kraussiana*.

S. dichroa, *Hort*. *See Vogelii*.

S. Douglasii, *Spring*. United States.

S. emiliana, *Hort*.

S. erythropus, *Spring*. Tropical America.

Var. *minor*, *Hort*.

S. filicina, *Spring*. *See hæmatodes*.

S. flabellata, *Spring*. Tropics.

S. flagellifera, *Hort*. *See plumosa*.

S. flexuosa, *Spring*. South Brazil.

Selaginella, *Spring*—*cont.*

S. Galeottei, *Spring*. Mexico.

S. grandis, *Moore*. Borneo.

S. Griffithii, *Spring*. Tropical Asia.

S. hæmatodes, *Spring*. Tropical America.

S. inæqualifolia, *Spring*. Tropical Asia.

Var. *bellula* (*Moore*). See var. *perelegans*.

Var. *perelegans* (*Moore*).

S. involvens, *Spring*. China and Japan.

Var. *texta*, *Baker*.

Var. *variegata*, *Hort*.

S. kraussiana, *A. Br.* South Africa, Azores, &c.

Var. *aurea*, *Hort*.

Var. *Brownii*, *Hort*.

Var. *Stansfieldii*, *Hort*.

Var. *variegata*, *Hort*.

S. lævigata, *Baker*. Madagascar.

Var. *Lyallii* (*Spring*).

S. lepidophylla, *Spring*. Tropical America.

S. leptostachya, *Bailey*. See *bakeriana*.

S. Lyallii, *Spring*. See *lævigata*.

S. Martensii, *Spring*. Mexico.

Var. *divaricata*, *Hort*.

Var. *stolonifera*, *A. Br.*

Var. *variegata*, *Hort*.

S. Mettenii, *A. Br.* Garden origin.

S. molliceps, *Spring*. Upper Guinea.

S. mutabilis, *Hort*. See *serpens*.

S. patula, *Spring*. West Indies.

S. pilifera, *A. Br.* Texas.

... of the ...
... of the ...

... of the ...
... of the ...

... of the ...
... of the ...

... of the ...
... of the ...

Selaginella, Spring—cont.

S. plumosa, *Baker*. Tropical Asia.

Var. *flagellifera*, *Hort*.

S. radiata, *Baker*. Tropical America.

S. radicata, *Spring*. See *plumosa*.

S. rubella, *Moore*. Garden origin.

S. rubricaulis, *Hort*. See *suberosa*.

S. serpens, *Spring*. West Indies.

S. stenophylla, *A. Br.* Mexico.

Var. *albospecta*, *Hort*.

S. suberosa, *Spring*. Tropical Asia.

S. texta, *Hort*. See *involuta*.

S. uncinata, *Spring*. China.

S. usta, *Vieill.* New Caledonia.

S. Victoriae, *Moore*. Borneo.

S. viridangula, *Spring*. Polynesia.

S. viticulosa, *Klotzsch*. Tropical America.

S. Vogelii, *Spring*. Tropical Africa.

Var. *dichroa* (*Hort.*).

S. Wallichii, *Spring*. Tropical Asia.

S. Warszewiczii, *Klotzsch*. See *radiata*.

S. Willdenovii, *Baker*. Tropical Asia.

Tmesipteris, Bernh.

T. Forsteri, *Endl.* See *tannense*.

T. tannense, *Bernh.* Australia and Polynesia.

III. APPENDIX.

Cultural forms of British Ferns.

Adiantum, *L.**A. Capillus-veneris*, *L.*

- Var. *admirable*, *Lowe*.
- Var. *concinnum*, *Hort*.
- Var. *cornubiense*, *Moore*.
- Var. *daphnites*, *Lowe*.
- Var. *fissum*, *Moore*.
- Var. *grande*, *Moore*.
- Var. *imbricatum*, *Lowe*.
- Var. *laciniatum*, *Hort*.
- Var. *magnificum*, *Fraser*.
- Var. *Mariesii*, *Moore*.
- Var. *maximum*, *Moore*.
- Var. *obrienianum*, *Birkenhead*.
- Var. *optandrum*, *Lowe*.
- Var. *perfectum*, *Padley*.
- Var. *Roperi*, *Hort*.
- Var. *semilunatum*, *Lowe*.
- Var. *tenuis*, *Hort*.
- Var. *undulatum*, *Moore*.

Aspidium, *Swartz* (*Polystichum*).*A. aculeatum*, *Sw.* Europe.

- Var. *acrocladon*, *Lowe*.
- Var. *Barnesii*, *Jones*.

Aspidium, Swartz (Polystichum)—cont.**A. aculeatum, Sw. Europe—cont.**Var. *Braunii*, *Hort.*Var. *capitatum*, *Jones.*Var. *cristatum*, *Moore.*Var. *cruciatum*, *Jones.*Var. „ *polydactylon.*Var. *dubium*, *Woll.*Var. *grandiceps*, *Jones.*Var. *Kitsonae*, *Jones.*Var. *meritum*, *Lowe.*Var. *multifidum*, *Woll.*Var. *proliferum*, *Woll.*Var. *pulcherrimum*, *Jones.*Var. *subcristatum*, *Jones.***A. angulare,* Willd. Europe.**Var. *acuto-dissectum*, *Carbonell.*Var. *acuto-divaricatum*, *Carbonell.*Var. *acuto-divisilobum*, *Carbonell.*Var. *acutilobum*, *Woll.*Var. „ *acutum*, *Lowe.*Var. „ *proliferum*, *Moore.*Var. *alatum*, *Moore.*Var. „ *pterophorum*, *Moore.*Var. *apuaeforme*, *Moore.*Var. *assimile*, *Jones.*Var. *athyroides*, *Woll.*Var. *Bayliae*, *Lyell.*Var. *brachiatum*, *Moore.*Var. „ *cristatum*, *Jarvis.*

* This is a sub-species of *A. aculeatum*, but for garden purposes it is more convenient to treat it as a species.

Aspidium, Swartz (Polystichum)—cont.**A. angulare, Willd. Europe—cont.**

Var. Buckleae, *Lowe*.

Var. calomelanos, *Jones*.

Var. capitatum, *Clap*.

Var. caudatum, *Carbonell*.

Var. Claphami, *Jones*.

Var. concinnum, *Moore*.

Var. confluens, *Moore*.

Var. congestum, *Moore*.

Var. conspicuulobum, *Lowe*.

Var. „ acutum, *Carbonell*.

Var. cristatum, *Moore*.

Var. „ biceps, *Carbonell*.

Var. „ gracile, *Woll*.

Var. cruciato-polydactylum, *Jones*.

Var. cruciatum, *Lowe*.

Var. „ majus, *Jones*.

Var. „ proliferum, *Hort*.

Var. curvato-pinnum, *Carbonell*.

Var. decompositum, *Moore*.

Var. „ densum, *Carbonell*.

Var. „ grande, *Jones*.

Var. „ grandiceps, *Jones*.

Var. „ plumosum, *Jones*.

Var. „ polydactylum, *Jones*.

Var. „ robustum, *Jones*.

Var. „ splendens, *Jones*.

Var. decurrens, *Moore*.

Var. densum, *Moore*.

Var. „ polydactylum, *Jones*.

Var. divisilobum, *Woll*.

Aspidium, *Swartz* (*Polystichum*)—*cont.*A. *angulare*, *Willd.* Europe—*cont.*Var. *divisilobum acutum*, *Jones.*Var. „ *cristatum*, *Ivery.*Var. „ *decorum*, *Jones.*Var. „ *deltoideum*, *Jones.*Var. „ *densum*, *Carbonell.*Var. „ *elegans*, *Jones.*Var. „ *foliosum*, *Jones.*Var. „ *grandiceps*, *Jones.*Var. „ *laxum*, *Jones.*Var. „ *lineare*, *Jones.*Var. „ *multifidum*, *Jones.*Var. „ *polydactylum*, *Jones.*Var. „ *proliferum*, *Woll.*Var. „ *reflexum*, *Jones.*Var. „ *robustum*, *Jones.*Var. „ *splendidissimum*, *Jones.*Var. *flexuosum*, *Woll.*Var. „ *polydactylum*, *Jones.*Var. *folioso-decompositum*, *Jones.*Var. *foliosum*, *Woll.*Var. „ *bulbiferum*, *Lowe.*Var. *frondosum*, *Jones.*Var. „ *decompositum*, *Jones.*Var. „ *polydactylum*, *Jones.*Var. *gracile*, *Woll.*Var. *grandiceps*, *Moore.*Var. *grandidens*, *Jones.*Var. „ *minus*, *Sim.*Var. „ *truncatum*, *Woll.*Var. *imbricatum*, *Moore.*Var. „ *congestum*, *Jones.*

Aspidium, Swartz (Polystichum)—cont.**A. angulare, Willd. Europe—cont.**Var. *inciso-acutum, Jones.*Var. *inciso-laciniatum, Jones.*Var. *incisum, Woll.*Var. „ *variegatum, Woll.*Var. *interruptum, Woll.*Var. *Kitsonae, Moore.*Var. *laciniatum, Woll.*Var. *lanceolatum, Moore.*Var. *latifolium, Woll.*Var. „ *grandiceps, Jones.*Var. *laxo-crispum, Jones.*Var. *lineare, Moore.*Var. „ *minus, Jones.*Var. „ *polydactylum, Jones.*Var. *lineatum, Moly.*Var. *multifidum, Woll.*Var. „ *densum, Jones.*Var. *multilobum, Woll.*Var. „ *deltoideum, Jones.*Var. „ *densum, Jones.*Var. „ *latum, Jones.*Var. „ *laxum, Jones.*Var. „ *multifidum, Jones.*Var. „ *ovale, Woll.*Var. „ *polydactylum, Jones.*Var. „ *revolvens, Jones.*Var. *oxyphyllum, Moore.*Var. *parvissimum, Moore.*Var. *pendens, Jones.*Var. *perserratum, Jones.*

Aspidium, Swartz (*Polystichum*)—*cont.**A. angulare*, Willd. Europe—*cont.*

Var. *plumosum*, Woll.

Var. „ *elegantissimum*, Jones.

Var. *polydactylum*, Moore.

Var. „ *capitatum*, Jones.

Var. „ *cruciatum*, Lowe.

Var. „ *grande*, Jones.

Var. „ *grandiceps*, Jones.

Var. „ *laxum*, Jones.

Var. „ *multifidum*, Jones.

Var. „ „ *grandiceps*, Jones.

Var. „ *robustum*, Jones.

Var. *proliferum*, Moore.

Var. „ *densum*, Carbonell.

Var. „ *holeanum*, Moore.

Var. *pulchellum*, Woll.

Var. *pulcherrimum*, Moore.

Var. *pulchrum*, Moore.

Var. *ramo-variens*, Jones.

Var. *remoto-decurrens*, Woll.

Var. *revolvens*, Moore.

Var. *revolvens grandiceps*, Jones.

Var. „ *polydactylum*, Jones.

Var. *scalptum*, Carbonell.

Var. *Stansfieldii*, Lowe.

Var. *stipatum*, Woll.

Var. *stipitatum*, Moore.

Var. *stipulatum*, Jones.

Var. *subplumosum*, Woll.

Var. *Thompsonae*, Moore.

Var. *torto-cristatum*, Padley.

Aspidium, Swartz (Polystichum)—cont.**A. angulare, Willd. Europe—cont.**Var. tripinnatum, *Moore*.Var. „ acutum, *Moore*.Var. „ densum, *Jones*.Var. „ laciniatum, *Jones*.Var. „ polydactylum, *Jones*.Var. venustum, *Moore*.Var. Widworthyi, *Carbonell*.Var. Wollastoni, *Lowe*.Var. „ majus, *Hort*.**A. Lonchitis, Sw. North temperate zone.****Asplenium, L.****A. Adiantum-nigrum, L. Europe, &c.**Var. acutum, *Pollini*.**A. Ceterach, L. Europe.**Var. major, *O'Kelly*.**A.^s Filix-fœmina, Bernh. Europe.**Var. abasiphyllum, *Lowe*.Var. acrocladon, *Moore*.Var. „ densum, *Stansf*.Var. adpresso-cristatum, *Jones*.Var. angustatum, *Lowe*.Var. apuæforme, *Barnes*.Var. axminsterense, *Lowe*.Var. canaliculatum, *Stansf*.Var. cancellatum, *Sim*.Var. „ cristatum, *Jones*.Var. ceratophyllum, *Mapp*.Var. clarissima, *Jones*.Var. caudiculatum, *Mapp*.

Asplenium, *L.*—*cont.*A. *Filix-fœmina*, *Bernh.* Europe—*cont.*

- Var. *congestum-cristatum*, *Stansf.*
- Var. *congestum findlayanum*, *Stansf.*
- Var. „ *grandiceps*, *Stansf.*
- Var. *conoides*, *Appleby.*
- Var. „ *cristatum*, *Stansf.*
- Var. *cornutum*, *Carbonell.*
- Var. *coronatum*, *Lowe.*
- Var. *corymbiferum*, *Moore.*
- Var. *Craigii*, *Moore.*
- Var. „ *angustatum*, *Jones.*
- Var. „ *laciniatum*, *Jones.*
- Var. *crispato-laciniatum*, *Jones.*
- Var. *crispum*, *Moore.*
- Var. „ *coronatum*, *Jones.*
- Var. „ *cristatum*, *Carbonell.*
- Var. *cristatum*, *Woll.*
- Var. „ *capitatum*, *Carbonell.*
- Var. „ *caudatum*, *Lowe.*
- Var. „ *corymbiferum*, *Carbonell.*
- Var. „ *defecto-furcillans*, *Mapp.*
- Var. „ *densum*, *Carbonell.*
- Var. „ *magnificum*, *Lowe.*
- Var. „ *reflexum*, *Carbonell.*
- Var. *cruciato-capitatum*, *Carbonell.*
- Var. „ *-coronatum*, *Carbonell.*
- Var. „ *-cristatum*, *Jones.*
- Var. „ *-gracile*, *Jones.*
- Var. „ *-grandiceps*, *Jones.*
- Var. „ *-magnificum*, *Jones.*
- Var. „ *-multifidum*, *Stansf.*

Asplenium, *L.—cont.*A. Filix-fœmina, *Bernh.* Europe—*cont.*

Var. cruciatum, *Jones.*

Var. „ angustum, *Lowe.*

Var. curtum, *Woll.*

Var. „ cristatum, *Stansf.*

Var. daphnites, *Glave.*

Var. „ capitatum, *Glave.*

Var. defecto-gracile, *Mapp.*

Var. diffidum, *Jones.*

Var. diffiso-multifidum, *Moore.*

Var. digitale, *Jones.*

Var. „ capitatum, *Jones.*

Var. „ purpureum, *Jones.*

Var. Elworthii, *Moore.*

Var. Fieldæ, *Moore.*

Var. „ minus, *Jones.*

Var. fissidens, *Moore.*

Var. „ elegans, *Mapp.*

Var. flabellatum, *Jones.*

Var. flabelliceps, *Carbonell.*

Var. flexuosum, *Woll.*

Var. foliosa-crispatum, *Carbonell.*

Var. „ -cristatum, *Woll.*

Var. Frizellæ, *Moore.*

Var. „ cristatum, *Jones.*

Var. „ multifidum, *Barnes.*

Var. „ nanum, *Ivery.*

Var. furcans, *Moore.*

Var. Girdlestoni, *Ivery.*

Var. glomeratum, *Moore.*

Var. „ grandiceps, *Jones.*

Asplenium, *L.*—*cont.*

A. *Filix-fœmina*, *Bernh.* Europe—*cont.*

- Var. *gracile*, *Moore*.
- Var. *grammicon*, *Lowe*.
- Var. *grandiceps*, *Moore*.
- Var. „ *globosum*, *Sim*.
- Var. „ *minus*, *Carbonell*.
- Var. „ *pumilum*, *Sim*.
- Var. *Harrisæ*, *Lowe*.
- Var. *Howardæ*, *Moore*.
- Var. *imbricatum*, *Carbonell*.
- Var. *incisum*, *Newm*.
- Var. *inæquale*, *Woll*.
- Var. „ *cristatum*, *Jones*.
- Var. *interruptum*, *Moore*.
- Var. *irregulare*, *Moore*.
- Var. *kalothrix*, *Lowe*.
- Var. *kilmorensis*, *Carbonell*.
- Var. *laceratum*, *Moore*.
- Var. *laciniatum*, *Moore*.
- Var. „ *angustatum*, *Jones*.
- Var. „ *congestum*, *Stansf*.
- Var. „ *Craigii*, *Jones*.
- Var. „ *cristatum*, *Lowe*.
- Var. „ *elegans*, *Stansf*.
- Var. „ *flabelliforme*, *Jones*.
- Var. „ *furcans*, *Jones*.
- Var. „ *ramulosum*, *Stansf*.
- Var. *Lowæ*, *Lowe*.
- Var. „ *capitatum*, *Lowe*.
- Var. *minimum*, *Birkenhead*.
- Var. *Moorei*, *Jones*.

*Asplenium, L.—cont.***A. Filix-fœmina, Bernh.** Europe—*cont.*Var. *multiceps, Moore.*Var. *multifidum, Moore.*Var. „ *cristatum, Jones.*Var. „ *laxum, Jones.*Var. „ *tenue, Jones.*Var. *nodosum, Jones.*Var. „ *multifidum, Jones.*Var. „ *nanum, Jones.*Var. „ *ramulosum, Jones.*Var. „ *truncatum, Jones.*Var. *orbiculatum, Lowe.*Var. *pannosum, Moore.*Var. „ *cristatum, Jones.*Var. *Parsonsi, Moore.*Var. *paucidentatum, Mapp.*Var. „ *cristatum, Mapp.*Var. *pectinatum, Moore.*Var. „ *Stableri, Mapp.*Var. *percristatum, Lowe.*Var. *perfectum, Carbonell.*Var. *phylomanes, Mapp.*Var. *plumosum, Moore.*Var. „ *divaricatum, Moore.*Var. „ *Edwardsi, Sim.*Var. „ *elegans, Parsons.*Var. „ *Jacksoni, Carbonell.*Var. „ *kalothrix, Lowe.*Var. „ *multifidum, Stansf.*Var. *polydactylum, Moore.*Var. „ *grandiceps, Jones.*Var. *Pritchardii, Stansf.*

*Asplenium, L.—cont.***A. Filix-fœmina, Bernh. Europe—cont.**

Var. „ *cristatum, Stansf.*

Var. *proteoides, Lowe.*

Var. *proteum, Clap.*

Var. „ *cristatum, Jones.*

Var. *pulcherrimum, Jones.*

Var. „ *cristatum, Jones.*

Var. *Pulleri, Moore.*

Var. *ramosum ramo-cristatum, Jones.*

Var. *rectangulare, Moore.*

Var. *refractum, Moore.*

Var. *reflexum, Woll.*

Var. *reflexum cristatum, Jones.*

Var. *regale, Moore.*

Var. *rhæticum, Moore.*

Var. „ *reflexum, Mapp.*

Var. *rotundatum, Woll.*

Var. „ *lunulatum, Carbonell.*

Var. *Ryleyæ, Carbonell.*

Var. *sagittatum, Moore.*

Var. „ *lunulatum, Carbonell.*

Var. *setigerum, Jones.*

Var. *Stableri, Lowe.*

Var. *Stansfieldii, Lowe.*

Var. *stipatum, Moore.*

Var. „ *capitatum, Carbonell.*

Var. *subplumosum, Padley.*

Var. „ *cristatum, Carbonell.*

Var. *todeoides, Stansf.*

Var. *tortile, Moore.*

Var. *torto-cristatum, Carbonell.*

Asplenium, L.—cont.**A. Filix-fœmina, Bernh. Europe—cont.**

- Var. „ -grandiceps, *Carbonell*.
- Var. truncato-pinnulum, *Jones*.
- Var. unco-cristatum, *Jones*.
- Var. „ -ramulosum, *Stansf.*
- Var. variegatum, *Wol.*
- Var. *Vernonæ*, *Jervis*.
- Var. „ corymbiferum, *Stansf.*
- Var. „ crispum, *Stansf.*
- Var. „ cristatum, *Mapp.*
- Var. *Victoriæ*, *Lowe*.
- Var. *Williamsii*, *Jones*.

A. Trichomanes, L. Temperate regions.

- Var. cristatum, *Moore*.
- Var. incisum, *Moore*.
- Var. trogyense, *Lowe*.

C. fragilis, Bernh. Europe.

- Var. angustata, *Smith*.
- Var. dickieana, *Moore*.

Lomaria, Willd.**L. Spicant, Desv. Europe.**

- Var. apiculata, *Moore*.
- Var. caudata, *Moore*.
- Var. concinna, *Moore*.
- Var. congesta, *Barnes*.
- Var. lineare, *Moore*.
- Var. multifurcata, *Moore*.
- Var. plumosa, *Barnes*.

Account of the ...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

Lomaria, Willd.—cont.

L. Spicant, Desv. Europe—*cont.*

Var. ramo-cristata, *Woll.*

Var. trinervia, *Moore.*

Nephrodium, Rich. (Lastrea.)

N. æmulum, Baker. Europe.

Var. cristatum, *Jones.*

N. Filix-mas, Desv. Europe.

Var. attenuatum, *Padley.*

Var. Barnesii, *Moore.*

Var. Bollandæ, *Moore.*

Var. capitatum, *Carbonell.*

Var. Clowesii, *Moore.*

Var. crispatisimum, *Jones.*

Var. crispulatum, *Hort.*

Var. crispum, *Sim.*

Var. „ gracile, *Lyell.*

Var. crista-galli, *Carbonell.*

Var. cristatum, *Lowe.*

Var. „ angustatum, *Moore.*

Var. „ caudatum, *Ivery.*

Var. cronkleyense, *Lowe.*

Var. depauperatum, *Padley.*

Var. digitatum, *Lowe.*

Var. Ellacombei, *Lowe.*

Var. elongatum, *Moore.*

Var. erosum, *Clowes.*

Var. „ crispatum, *Carbonell.*

Var. fluctuosum, *Moore.*

Var. furcans, *Moore.*

Var. grandiceps, *Sim.*

Var. Ingrami, *Moore.*

Nephrodium, Rich. (Lastrea)—cont.

N. Filix-mas, Desv. Europe—cont.

- Var. interruptum, *Moore*.
- Var. intermedium, *Carbonell*.
- Var. Jervisii, *Moore*.
- Var. lineare, *Woll*.
- Var. Marsdeniæ, *Lowe*.
- Var. multiforme, *Barnes*.
- Var. paleaceum, *Don*. (pseudo-mas, *Woll*.)
- Var. „ Belperi, *Lowe*.
- Var. „ cristatum, *Moore*.
- Var. „ „ intermedium, *Sim*.
- Var. „ furcillatum, *Woll*.
- Var. „ Lyelli, *Lowe*.
- Var. „ ramo-furcillatum, *Woll*.
- Var. „ recurvum, *Moore*.
- Var. „ revolvens, *Woll*.
- Var. polydactylum, *Lowe*.
- Var. propinquum, *Woll*.
- Var. „ cristatum, *Moore*.
- Var. „ excurrens, *Moore*.
- Var. „ Gottii, *Lowe*.
- Var. „ gracile, *Moore*.
- Var. „ grandiceps, *Moore*.
- Var. „ laciniatum, *Lowe*.
- Var. „ parviceps, *Barnes*.
- Var. „ polydactylum, *Barnes*.
- Var. Pulleri, *Jones*.
- Var. ramo-cristatum, *Stansf*.
- Var. „ -furcillatum, *Carbonell*.
- Var. ramosum, *Barnes*.
- Var. ramulosissimum, *Woll*.

Nephrodium, Rich. (Lastrea)—cont.**N. Filix-mas, Desv. Europe—cont.**Var. *revolvens*, *Humphreys*.Var. *subcristatum*, *Carbonell*.Var. *variegatum*, *Lowe*.**N. montanum, Baker. Europe.**Var. *cristatum*, *Woll*.**N. spinulosum, Desv. Europe.**Var. *dilatatum* (*Desv.*).Var. „ *angusti-pinnulum*, *Sim*.Var. „ *calomelanos*, *Jones*.Var. „ *crispato-cristatum*, *Jones*.Var. „ *crispum*, *Woll*.Var. „ *grandiceps*, *Barnes*.Var. „ *Howardæ*, *Jones*.Var. *lepidotum*, *Moore*.**Osmunda, L.****O. regalis, L. Europe, &c.**Var. *cristata*, *Moore*.Var. *palustris*, *Hort*.Var. *purpurascens*, *Moore*.**Polypodium, L.****P. alpestre, Hoppe. Europe.**Var. *flexile*, *Newm*. Scotland.**P. Phegopteris, L. Europe.**Var. *cristatum*, *Jones*.Var. *strictum*, *Barnes*.**P. vulgare, L. Europe.**Var. *acuto-serratum*, *Jones*.Var. *attenuatum*, *Moore*.Var. *bifidum*, *Francis*.

P. vulgare, L. Europe—*cont.*

- Var. bifido-cristatum, *Mapp.*
- Var. cambricum (*L.*).
- Var. „ Barrowi, *Barnes.*
- Var. cornubiense, *Moore.*
- Var. „ elegantissimum, *Stansf.*
- Var. „ Fowleri, *Lowe.*
- Var. cornutum, *Jones.*
- Var. crenatum, *Moore.*
- Var. crispum, *Jones.*
- Var. cristatum, *Moore.*
- Var. „ Morleyi, *Jones.*
- Var. dentatum, *Moore.*
- Var. dentatum Barnesii, *Moore.*
- Var. „ Whitakeri, *Moore*
- Var. divisilobum, *Hort.*
- Var. Elworthii, *Ivery.*
- Var. falcatum, *Jones.*
- Var. folioso-abruptum, *Barnes.*
- Var. glomeratum, *Moly.*
- Var. irregulare, *Moore.*
- Var. marginatum, *Moore.*
- Var. marginato-attenuatum, *Jones.*
- Var. multifido-cristatum, *Stansf.*
- Var. omnilacerum, *Moore.*
- Var. „ truncatum, *Wilson.*
- Var. pulcherrimum, *Moore.*
- Var. ramosum, *Moore.*
- Var. „ Hillmani, *Carbonell.*
- Var. semilacerum (*Link.*).
- Var. „ obtusum, *Dadds.*
- Var. „ truncatum, *Barnes.*
- Var. serratum, *Moore.*

Polypodium, L.—cont.**P. vulgare, L.** Europe—*cont.*Var. *suprasoriferum, Woll.*Var. *trichomanoides, Backh.*Var. *truncatum, Lowe.***Pteris, L.****P. aquilina, L.** Cosmopolitan.Var. *cristata, Moore.***Scolopendrium, Smith.****S. vulgare, L.** Europe.Var. *aduncum, Moore.*Var. *angustifrons, Ivery.*Var. *bimarginatum, Moore.*Var. *bimarginato-tenue, Jones.*Var. *bulbiferum, Lowe.*Var. *chelæfrons, Woll.*Var. *concevum, Lowe.*Var. *Coolingii, Lowe.*Var. *cornutum, Lowe.*Var. *corrugatum, Lowe.*Var. *crispum, Grey.*Var. „ *Cowburni, Lowe.*Var. „ *densum, Sim.*Var. „ *fimbriatum, Lowe.*Var. „ *grande, Lowe.*Var. „ *latum, Moore.*Var. „ *majus, Moore.*Var. „ *multifidum, Jones.*Var. „ *reflexum, Wills.*Var. „ *robustum, Jones.*

Scolopendrium, *Smith*—*cont.*

S. vulgare, *L.* Europe—*cont.*

Var. *crispum*, *serratum*, *Jones*.

Var. *crista-galli*, *Woll*.

Var. *cristatum*, *Moore*.

Var. „ *variegatum*, *Carbonell*.

Var. *cristatum cristato-digitatum*, *Sim*.

Var. *cuspidis*, *Carbonell*.

Var. *elegantissimum*, *Hort*.

Var. *fissidens*, *Woll*.

Var. *fisso-crenatum*, *Jones*.

Var. *fissum*, *Moore*.

Var. *Fowleri*, *Hort*.

Var. *grandiceps*, *Jones*.

Var. *Hankeyi*, *Lowe*.

Var. *illustre*, *Lowe*.

Var. *involvens*, *Mapp*.

Var. „ *minus*, *Hort*.

Var. *iveryanum*, *Ivery*.

Var. *Kelwayi*, *Lowe*.

Var. *laceratum*, *Moore*.

Var. *labyrinthum*, *Lowe*.

Var. *lonchophorum*, *Moore*.

Var. *macrosorum*, *Moore*.

Var. *Moonæ*, *Lowe*.

Var. *Malcolmsonæ*, *Stansf*.

Var. *marginale*, *Hort*.

Var. *marginatum*, *Moore*.

Var. „ *fissum*, *Lowe*.

Var. *marginato-angustum*, *Clap*.

Var „ *-multifidum*, *Moore*.

Scolopendrium, *Smith*—*cont.*S. vulgare, *L.* Europe—*cont.*

Var. marginato-muricatum, *Jones.*

Var. muricatum, *Lowe.*

Var. muricato-cristatum, *Woll.*

Var. omnilacerum, *Lowe.*

Var. „ truncatum, *Hort.*

Var. peraferens, *Woll.*

Var. pulcherrimum, *Hort.*

Var. ramoso-minus, *Jackson.*

Var. ramulosissimum, *Woll.*

Var. scalpturatum, *Moore.*

Var. submarginato-fissum, *Wills.*

Var. supralineatum, *Moore.*

Var. undulatum, *Moore.*

Var. „ capitatum, *Hort.*

Var. „ cochleatum, *Jones.*

Var. „ marginatum, *Jones.*

Var. „ reflexum, *Hort.*

Var. *Veloisii*, *Moore.*

Var. variable, *Lowe.*

Var. variegatum, *Woll.*

Var. „ aureolum, *Lowe.*

GUIDES ON SALE AT THE ROYAL GARDENS, KEW.

Key Plan and Index to the Royal Botanic Garden and Arboretum, Kew. 3rd ed. 1893. *Price 2d.*

Official Guide to the Museums of Economic Botany. No. 1: Dicotyledons and Gymnosperms. 2nd ed., revised and augmented. 1886. *Price 4d.*

Official Guide to the Museums of Economic Botany. No. 2: Monocotyledons and Cryptogams. 1894. *Price 4d.*

Official Guide to the Museums of Economic Botany. No. 3: Timbers. 2nd ed., revised and augmented. 1893. *Price 3d.*

Official Guide to the North Gallery. 5th ed., revised and augmented. 1892. *Price 6d.*

Hand-list of Trees and Shrubs grown in Arboretum. Part I.: Polypetalæ. 1894. *Price 8d.*

Hand-list of Ferns and Fern-Allies cultivated in the Royal Gardens. 1895. *Price 6d.*
